

Scaling laws and effective dimension in lattice $SU(2)$ Yang-Mills theory with a compactified extra dimension

Shinji Ejiri

Department of Physics, University of Wales Swansea, Swansea, SA2 8PP, United Kingdom

Shouji Fujimoto and Jisuke Kubo

Institute for Theoretical Physics, Kanazawa University, Kanazawa 920-1192, Japan

(Received 24 April 2002; published 14 August 2002)

Monte Carlo simulations are performed in a five-dimensional lattice $SU(2)$ Yang-Mills theory with a compactified extra dimension, and scaling laws are studied. Our simulations indicate that, as the compactification radius R decreases, the confining phase spreads more and more to the weak coupling regime, and the effective dimension of the theory changes gradually from five to four. Our simulations also indicate that the limit $a_4 \rightarrow 0$ with R/a_4 kept fixed exists in both the confining and deconfining phases if R/a_4 is small enough, where a_4 is the lattice spacing in the four-dimensional direction. We argue that there exists a maximal radius above which the color degrees of freedom are not confined. Comments on deconstructing extra dimensions are given.

DOI: 10.1103/PhysRevD.66.036002

PACS number(s): 11.10.Kk, 11.10.Wx, 11.15.Ha, 11.25.Mj

I. INTRODUCTION

Since Kaluza and Klein [1] found that the electromagnetic and gravitational forces can be unified by introducing a fifth dimension, their idea has attracted attention for many decades. Recently, there has been a lot of renewed interest in field theories with extra dimensions, in which the length scale of the extra dimensions can be so large that they could be experimentally observed [2–4]. It is assumed that, for distances larger than the compactification size, the massive Kaluza-Klein excitations decouple so that these theories behave as a four-dimensional continuum theory at low energies. Since, however, Yang-Mills theories in more than four dimensions are nonrenormalizable, it is not at all clear that the infinite tower of the Kaluza-Klein excitations decouples even if each massive excitation is suppressed: A naive expectation of their contribution would be $\infty \cdot 0$.

In four dimensions, the color degrees of freedom are confined even for a weak gauge coupling. How can a confining four-dimensional Yang-Mills theory emerge from a higher-dimensional Yang-Mills theory which is deconfining in the weak coupling regime [5–7]? Although the assumption of the decoupling of the Kaluza-Klein excitations sounds physically correct, it is by no means trivial to show that they nonperturbatively decouple in such a way that the color confinement takes place even at weak gauge coupling. Recently, we [8] started to address related problems in a concrete example, namely, the pure lattice $SU(2)$ Yang-Mills theory in five dimensions with one dimension compactified on a circle. We observed [8] that the compactification changes the nature of the phase transition, and that a second order phase transition, which does not exist in the uncompactified case, occurs, thus confirming the long-standing expectation of Lang, Pilch, and Skagerstam [6]. The phase is defined by the Polyakov loop that extends into the fifth dimension, and the phase transition is expected to be of second order, because the compactified $SU(2)$ lattice gauge theory in five dimensions belongs to the universality class of the Z_2 spin model in four

dimensions. For the first time, we [8] computed the lattice β function in a Yang-Mills theory in more than four dimensions, and verified nonperturbatively the power-law running of the gauge coupling constant [4,9–11].

In this paper we would like to extend the analyses of [8]. We first observe that if the compactification radius becomes smaller and smaller, the confining phase spreads more and more to the weak coupling regime. At the same time we compute the effective dimension [12,13], and see that the theory behaves more and more as a four-dimensional Yang-Mills theory. Based on this result, we argue that the color degrees of freedom are confined only for $R < R_{\text{max}}$. The confining phase is defined by the string tension between two static quarks that are separated in the four-dimensional subspace. This definition of phase should not be confused with the definition by the Polyakov loop that extends into the fifth dimension, which was mentioned above.

Our calculations of the potential between two static quarks separated in the four-dimensional subspace show that the deconfining phase is a Coulomb phase. We then discuss the nature of the transition from the deconfining phase to the confining phase for fixed values of R/a_4 , where a_4 is the lattice spacing in the four-dimensional direction. We confirm that if R/a_4 is small enough, it is consistent with a second order transition. Combined with the result of [8], we therefore come to the conclusion that, as we decrease the value of RA , the first order transition for large values of RA changes to a crossover transition, and finally becomes of second order.

We give some nonperturbative comments on deconstructing extra dimensions [14] in the Conclusion.

II. EFFECTIVE DIMENSION

In order to take into account the compactification effects in this theory, it is crucial to use anisotropic lattices [15] which have different lattice spacings a_4 and a_5 in the four-dimensional directions and in the fifth direction. For definite-

ness we employ the Wilson action for pure $SU(2)$ lattice gauge theory:

$$S = \frac{\beta}{\gamma} \sum_{P_4} \left[1 - \frac{1}{2} \text{Re Tr } U_{P_4} \right] + \beta \gamma \sum_{P_5} \left[1 - \frac{1}{2} \text{Re Tr } U_{P_5} \right], \quad (2.1)$$

where U_{P_4} denote plaquette variables in the four-dimensional sublattice, and U_{P_5} are those that are extended into the fifth dimension. The gauge coupling constant g_5 has the dimension of $\sqrt{a_4}$, which is related to β by

$$a_4 g_5^{-2} = \beta/4 \quad (2.2)$$

at the tree level. Periodic boundary conditions are imposed in all directions, and we use a lattice size of the form $N_4^4 \times N_5$ (we mostly use $N_4=12$ and $N_5=4$). The compactification radius is defined as $R = a_5 N_5 / 2\pi$ if $N_4 a_4 > N_5 a_5$ is satisfied, and the correlation-anisotropy parameter is defined as $\xi = a_4 / a_5$. The tree level relation $\gamma = \xi$ will be modified at the quantum level [15], and throughout this paper we assume that the ξ - γ relation obtained in [8] is satisfied in both the confining and deconfining phases. Simulations are performed for

$$\gamma = 3.6, 4.0, 4.6, 5.0, \quad (2.3)$$

which is equivalent to [8]

$$\frac{2\pi R}{a_4} = \frac{N_5 a_5}{a_4} = \frac{4}{\xi} \approx 0.72, 0.64, 0.55, 0.50, \quad (2.4)$$

where we have used $N_5=4$ above. We chose this range of $2\pi R/a_4$ because we expect from the previous calculations [8] that the limit $2\pi R/a_4 \rightarrow 0$ may exist, and we observe some scaling behavior.

To define the physical scale in the confining phase, we use the string tension σ between two static quarks that are separated in the four-dimensional subspace. Since the string tension is a physical quantity, the lattice string tension σ_L should behave like a_4^2 as $a_4 \rightarrow 0$, where a_4 can be related by the β function to the dimensionless bare gauge coupling

$$g^2 = \frac{8\pi}{\beta} = \Lambda g_5^2, \quad (2.5)$$

where we have identified Λ with $2\pi/a_4$ because $4 \times (\pi/a_4)^2 = (2\pi/a_4)^2$. Since we expect that the massive Kaluza-Klein excitations will decouple increasingly as $2\pi R/a_4$ decreases, the lattice β function β_L cannot assume a purely five- or four-dimensional form. Instead, we expect a continuous change of its form. This is quantitatively expressed by the so-called effective dimension D_{eff} , which is a function of $2\pi R/a_4$ [13]. So we assume that β_L can be written as

$$\beta_L = -a_4 \frac{dg^2}{da_4} = [D_{\text{eff}}(2\pi R/a_4) - 4]g^2 - \frac{2b}{16\pi^2} g^4, \quad (2.6)$$




FIG. 1. $\sqrt{\chi_0} = \sqrt{\sigma_L}$ versus β at $\gamma=5.0$. The filled symbols are obtained from the Creutz ratio (3.1) and the open ones are obtained from the static potential (3.2).

where $b = 22/3 - 2/3 = 20/3$. Therefore, the evolution equation for g^2 can be easily integrated in the case that $2\pi R/a_4$ is kept fixed while a_4 changes. We obtain for this case

$$\sqrt{\sigma_L} \sim a_4 \sim \left(\frac{2b}{16\pi^2} \frac{1}{D_{\text{eff}} - 4} - \frac{\beta}{8\pi} \right)^{1/(D_{\text{eff}} - 4)}. \quad (2.7)$$

It is important to notice that as $D_{\text{eff}} \rightarrow 4$ we obtain the logarithmic form

$$g^{-2} = \beta/8\pi = (2b/16\pi^2) \ln a_4 + \text{const.} \quad (2.8)$$

That is, if we can show that the effective dimension D_{eff} in the confining phase varies from 5 to 4 as $2\pi R/a_4$ decreases from a larger value to a smaller value, we show the continuous decoupling of the Kaluza-Klein excitations, and the confining phase spreads more and more to the weak coupling regime as R decreases.

III. CONFINING PHASE

Now we come to the results of our Monte Carlo simulations on a $12^4 \times 4$ lattice. We use the Creutz ratio $\chi(I, J)$ obtained from the rectangular Wilson loops $W(I, J)$ with lengths of I and J in the four-dimensional subspace. We assume that the Creutz ratio takes the form

$$\chi(I, J) = \chi_0 - \chi_1 \left(\frac{1}{I(I-1)} + \frac{1}{J(J-1)} \right) + \chi_2 \left(\frac{1}{I(I-1)J(J-1)} \right), \quad (3.1)$$

and we identify χ_0 with the lattice string tension σ_L . We generated 2500 configurations for each simulation point after thermalization, and the Wilson loops were measured every five configurations for the calculation of a Creutz ratio. Errors were estimated by the jackknife method. The filled symbols in Fig. 1 are the result obtained from the Monte Carlo simulations with $\gamma=5.0$, where the vertical axis stands for




FIG. 2. The scaling behavior of $\sigma_L^{1/2}$ for different values of γ . The solid lines are drawn by using Eq. (2.7), where D_{eff} is taken from Table I. The data with a filled symbol are used for the fit.

$\sqrt{\chi_0} = \sqrt{\sigma_L}$ and the horizontal axis stands for β . We have also calculated σ_L from the static potential to make sure that σ_L obtained from the Creutz ratios is reliable.¹ The static potential we have assumed has the form

$$V(X) = C_0 - C_1 \frac{1}{X} + C_2 \left(\frac{1}{X} - \left[\frac{1}{X} \right] \right) + C_3 X, \quad (3.2)$$

where $[1/X]$ is the three-dimensional Coulomb potential on a lattice, and is given by

$$\left[\frac{1}{X} \right] = 4\pi \int_{-\pi}^{\pi} \frac{d^3 p}{(2\pi)^3} \frac{\exp\left\{ i \sum_i X_i \sin(p_i/2) \right\}}{\sum_{i=1}^4 \sin^2(p_i/2)}. \quad (3.3)$$

The open symbols in Fig. 1 correspond to the result obtained from the static potential. Comparing the two results in Fig. 1 we see that the lattice string tensions obtained from the Creutz ratios agree with those obtained from the static potential. We made the same comparison for different values of γ , and found the same result. So in the following analyses we use only the lattice string tensions from the Creutz ratio, because we have more data for this case and we do not want to mix data obtained by two different methods.

We see from Fig. 1 that above $\beta \gtrsim 3.0$ the square root of the lattice string tension $\sqrt{\sigma_L}$ first decreases linearly until $\beta \sim 3.3$, and then its slope becomes milder. The tail for large β is certainly due to the finite lattice size effects, but the change from the linear decrease of $\sqrt{\sigma_L}$ to a milder one around $\beta \sim 3.3$ may indicate that the theoretical expectation (2.7) is correct. Although it is in principle possible to check by increasing the lattice size how much finite lattice size effects may be contained in the tail of $\sqrt{\sigma_L}$, it is impossible to do this at the moment because of the limitations of the

¹We give more details of calculating the static potential in Sec. V when calculating the potential in the Coulomb phase.




FIG. 3. The effective dimension as a function of the number n of data points that are used for a fit. We increase n starting from 4 until the value of $(D_{\text{eff}} - 4)^{-1}$ becomes stabilized.

computing facility given to us. Below we sketch how we confirm Eq. (2.7) and compute D_{eff} .

The effective dimension can be obtained by fitting the function (2.7) to the data. To this end, we first choose four neighboring data points that lie around the middle of the data set for a given γ , and using these points, we fit the function (2.7) to obtain the effective dimension. (In the case of $\gamma = 5.0$, for instance, we use the data points at $\beta = 3.20, 3.22, 3.24$, and 3.26 .) Then we increase the number of the data points to be used by 2 by including the next neighboring data point on both sides. In doing so, we obtain the effective dimension and also χ^2 per degree of freedom (DOF) as a function of number n of the data points that are used for the fit. We repeat the same analysis for the different values of $2\pi R/a_4$ given in Eq. (2.4). The results are shown in Figs. 2 and 3 and in Table I. In Fig. 3, the vertical axis stand for $(D_{\text{eff}} - 4)^{-1}$ and the error bar is computed from χ^2/DOF . We see that as n increases the error bar decreases and the central values converge. The results are summarized in Table I, and we see that the effective dimension D_{eff} decreases gradually from 4.7057(55) to 4.5230(82) as γ increases from 3.6 to 5.0, which means as $2\pi R/a_4$ decreases from 0.72 to 0.5 [see Eq. (2.4)].

The β^* in Table I is the value at which σ_L and hence a_4 should vanish if the theoretical assumption (2.7) is correct and is extrapolated for larger values of β (see also Fig. 2). We emphasize that our results indicate that the limit $a_4 \rightarrow 0$ with R/a_4 kept fixed exists in the confining phase at finite β .

TABLE I. Effective dimension for different values of γ ($\Lambda = 2\pi/a_4$).

γ	$R\Lambda$	D_{eff}	$(\beta_{\text{min}} : \beta_{\text{max}})$	χ^2/DOF	β^*
3.6	0.72	4.7057(55)	(2.30 : 2.76)	0.525	3.007(23)
4.0	0.64	4.6456(54)	(2.50 : 2.96)	0.438	3.286(22)
4.6	0.55	4.5695(55)	(2.80 : 3.26)	0.778	3.726(36)
5.0	0.50	4.5230(82)	(3.00 : 3.46)	0.598	4.057(43)

IV. THE MAXIMAL RADIUS

The same analysis in real QCD in Sec. III would constrain the size of the compactification radius in QCD, which we would like to estimate without detailed calculations. Our estimate below is based on many assumptions that cannot be justified at present, and so the number we obtain should not be taken seriously. But it is worthwhile to do this to see what kinds of problems are involved if one would like to do a more reliable estimate.

To see that there exists a maximal radius for color confinement in the four-dimensional subspace, we recall the results obtained in the previous section and those from the next section:

$$g^2 \geq (\leq) (D_{\text{eff}} - 4)(16\pi^2/2b) \quad (4.1)$$

for the (de)confining phase. Therefore, for a given value of D_{eff} , there should exist a smallest value of g^2 for color confinement to occur, which is $\sim (D_{\text{eff}} - 4)(16\pi^2/2b)$. The question is how g^2 can be related to the gauge coupling g_{KK}^2 of Kaluza-Klein theory, the four-dimensional theory with a Kaluza-Klein tower. At the tree level, it is $g_{\text{KK}}^2 = g^2(2\pi R\Lambda)^{-1}$, but in higher orders this relation will receive quantum corrections, where we have used $\Lambda^2 = (\pi/a_4)^2 \times 4$. To answer the question, we first assume that $D_{\text{eff}}(R\Lambda) \rightarrow 4$ (5) as $R\Lambda \rightarrow 0$ (∞), and we consider a redefinition of g^2 according to [12,13]:

$$g_{\text{KK}}^2 = \eta^{-1}(R\Lambda)g^2, \quad \eta(t) = \exp \int_0^t \frac{dt'}{t'} [D_{\text{eff}}(t') - 4]. \quad (4.2)$$

Note that the β function of g_{KK}^2 becomes

$$\beta_{\text{KK}} = -(2b/16\pi^2)\eta(R\Lambda)g_{\text{KK}}^4. \quad (4.3)$$

Since the function $\eta(R\Lambda)$ becomes proportional to $R\Lambda$ as $R\Lambda \rightarrow \infty$,² the new gauge coupling describes a power-law behavior [4,9–11]. Furthermore, we see from Eq. (4.2) that g_{KK}^2 approaches g^2 as $R\Lambda$ approaches 0. Recalling now the assumption that D_{eff} approaches 4 as $R\Lambda$ approaches 0 and Eq. (2.8), we see that the renormalization group flow of the new gauge coupling g_{KK}^2 for small $R\Lambda$ takes exactly the same form as the one for the effective, four-dimensional theory without the Kaluza-Klein tower. Therefore, we assume that g_{KK}^2 is the gauge coupling of the four-dimensional theory with the Kaluza-Klein tower.

Now, suppose that QCD results from a five-dimensional QCD. As we have argued above, g^2 becomes g_{KK}^2 at low energies, and we then identify $2\pi/a_4$ with the physical scale Λ of the effective theory, rather than with the ultraviolet cutoff. Since $g_R^2(M_Z)/4\pi \approx 0.12$ in QCD and $b=7$, the con-

straint (4.1) can be converted to that of an effective dimension, i.e., $D_{\text{eff}}(RM_Z) \leq 4.13$. Therefore, if we know the function $D_{\text{eff}}(t)$ exactly, we can calculate the range of t for which the inequality (4.1) is satisfied. From the results given in Table I we find that the effective dimension as a function of t can be written as $D_{\text{eff}}(t) \approx 4 + t$. Assuming that this function can be used even for small t , we then obtain $R\Lambda \leq 0.13$, which would imply that $1/R \geq O(1)$. TeV should be satisfied for the color degrees of freedom in QCD to be confined.

There are various problems involved in our estimates above, apart from the main assumptions that Eq. (2.6) is correct in QCD, the effective dimension $D_{\text{eff}}(t)$ can be extrapolated for smaller values of t although we know it only for $0.5 \leq t \leq 0.72$, and the form $D_{\text{eff}}(t) \approx 4 + t$ remains the same in QCD. One is the identification $2\pi/a_4 \approx \Lambda$, and the other is $g(\Lambda) \approx g_R(\mu = 2\pi/a_4)$, where $g_R(\mu)$ is a renormalized gauge coupling with the renormalization scale μ in a certain renormalization scheme. The first one comes from the assumption that we are very close to a continuum theory that possesses a four-dimensional rotational invariance. The second one comes from the fact that Λ_{lattice} and $\Lambda_{\overline{\text{MS}}}$ are not very much different in QCD so that the value of the bare lattice gauge coupling $g(\Lambda)$ is approximately equal to that of the renormalized gauge coupling $g_R(\mu = \Lambda)$. In order to justify these assumptions and obtain more reliable relations among them, we first of all have to refine and extend the γ - ξ relation given in Eqs. (2.3) and (2.4), which were obtained only for $\beta < 1.8$ for the $SU(2)$ theory in [8]. More important is that, apart from the fact that we have to do the calculations in the case of $SU(3)$, we should consider the continuum limit with the compactification radius R kept fixed. This will be necessary to introduce a real physical scale and to relate the string tension to R .

Therefore, our estimate of R_{max} above should not be taken seriously. However, simulations on five-dimensional, compactified $SU(3)$ lattice gauge theory would go beyond the scope of the present paper, and we would like to leave this problem to future work. The crucial point is that there exists a maximal radius.

V. COULOMB PHASE

The confining phase shrinks as R decreases, which we have already seen above. Next we would like to show that the deconfining phase is a Coulomb phase. To begin with, we consider the Wilson loop $W(\vec{x}, t)$ at the tree level in continuum perturbation theory. The static potential can be obtained from

$$\begin{aligned} V(x) &= \lim_{t \rightarrow \infty} [\ln W(\vec{x}, t)]/t \\ &= -\frac{3}{4}g_s^2 \frac{1}{2\pi R} \frac{1}{4\pi x} \coth\left(\frac{x}{2R}\right) \\ &= -\frac{3}{4}g_s^2 \times \begin{cases} \frac{1}{4\pi^2 x^2} & \left(\frac{x}{2R} \ll 1\right), \\ \frac{1}{2\pi R} \frac{1}{4\pi x} & \left(\frac{x}{2R} \gg 1\right). \end{cases} \end{aligned} \quad (5.1)$$

²The proportionality constant depends on D_{eff} as a function of t , which, however, depends on the regularization used [13]. Therefore, the lattice regularization does not reproduce the same coefficient [8] obtained in [4].




FIG. 4. The Coulomb potential (5.2). The filled symbols are the raw data points, and the dotted line is $V(X)$ of Eq. (5.5) with $C_0 = 0.3230(14)$ and $C_1 = 0.1086(30)$. The open symbols stand for the rotationally invariant data points.

We have the usual Coulomb potential for $x/2R \gg 1$, and we see that the dimensionless gauge coupling \hat{g} , normalized for four-dimensional Yang-Mills theory at the tree level, is given by $\hat{g} = g_5 / \sqrt{2\pi R}$, as is well known [3,4]. The corresponding expression on a lattice is

$$V_L(X) = \lim_{T \rightarrow \infty} \ln W(X, T) / W(X, T+1), \quad (5.2)$$

where $W(X, T)$ is a lattice Wilson loop. The lattice distances X and T are made dimensionless by dividing by a_4 . We are interested in the potential between two static quarks that are separated in four dimensions, and therefore X and T are supposed to be in the four-dimensional sublattice. Since in the actual calculations we cannot take the limit $T \rightarrow \infty$, we consider also off-axis loops and use the standard smearing techniques [16] to improve the convergence of approximants with increasing T . Our smearing procedure consists of iteratively replacing each spatial (three-dimensional) link by the sum of itself and the neighboring four spatial staples with a weight parameter ϵ :

$$\begin{aligned} U_i(x, y) &\rightarrow U_i^*(x, y) \\ &= \mathcal{P}_{SU(2)} \left(U_i(x, y) + \epsilon \sum_{j(i \neq j)=1}^3 F_{ij}(x, y) \right), \quad (5.3) \\ F_{ij}(x, y) &= U_j(x, y) U_i(x + \hat{j}, y) U_j^\dagger(x + \hat{i}, y) \\ &\quad + U_j^\dagger(x - \hat{i}, y) U_i(x - \hat{i}, y) U_j(x - \hat{i} + \hat{j}, y), \end{aligned}$$

where $\mathcal{P}_{SU(2)}$ denotes a projection operator back onto the $SU(2)$ manifold.

We generated 10000 configurations for each simulation point after thermalization, and the smeared Wilson loops were measured every 100 configurations for the calculation of the static potential. We iterated Eq. (5.3) 60 times with $\epsilon = 0.1$ in the case of the confining phase, 100 times with $\epsilon = 0.2$ in the case of the Coulomb phase. In Fig. 4 we show

the result (filled symbols) for the lattice potential $V_L(X)$ as a function of X at $\beta = 5.0$ and $\gamma = 5.0$ (which is equivalent to $2\pi R/a_4 = 0.5$). The condition $x/2R \gg 1$ to obtain a $1/X$ potential becomes $X \gg 1/2\pi$ in this case, and we assume that the lattice potential $V_L(X)$ takes the form

$$V_L(X) = C_0 - C_1 \frac{1}{X} + C_2 \left(\frac{1}{X} - \left[\frac{1}{X} \right] \right), \quad (5.4)$$

where $[1/X]$ (the three-dimensional Coulomb potential on a lattice) is given in Eq. (3.3). The first term of Eq. (5.4) is the unphysical self-energy, the second term is the rotationally invariant part of the Coulomb potential, and the third term is the most dominant part of its breaking. From a χ^2 fit we find that $C_0 = 0.3230(14)$, $C_1 = 0.1086(30)$, and $C_2 = 0.0776(27)$. The fitted lattice potential with the C_2 term in Eq. (5.4) suppressed, i.e.,

$$V(X) = C_0 - C_1 \frac{1}{X}, \quad (5.5)$$

is the dotted curve in Fig. 4, while the open symbols stand for the rotationally invariant data points. We see that the data justify the assumption that the deconfining phase is a Coulomb phase.

VI. NATURE OF THE PHASE TRANSITION

As the next task we consider the nature of the transition from the confining phase to the Coulomb phase. In the confining phase our data indicate that the limit $a_4 \rightarrow 0$ with R/a_4 kept fixed exists at finite β . If we can show that a_4 also vanishes at the same value of β in the Coulomb phase, the transition from the confining phase to the Coulomb phase is of second order.

To this end, we have to define the scale in the Coulomb phase. In the naive continuum theory there are two dimensional quantities, the gauge coupling g_5 and the compactification radius R . Therefore, we assume that R and the low-energy value of g_5 are independent physical quantities at the quantum level, too. We then consider the limit $a_4 \rightarrow 0$ with $2\pi R/a_4$ kept fixed, which is the same limiting process we have considered in the confining phase. In this limit, the quantity $g_5^2/2\pi R$ [the coefficient C_1 of the tree level Coulomb potential (5.4)] has to diverge because $R \rightarrow 0$ while g_5 should remain finite by assumption. So naively one expects the scaling law $C_1^{-1} \sim R \sim a_4 \sim (\beta - \beta^*)$, where β^* is the critical value of β at which $\sigma_L^{1/2} \sim a_4$ vanishes. In Fig. 5 we plot C_1^{-1} versus β for different values of γ [or $2\pi R/a_4$ of Eq. (2.4)]. We see that C_1^{-1} linearly decreases, and make therefore a theoretical ansatz for the scaling law:

$$C_1^{-1} = D_0 - D_1 \beta. \quad (6.1)$$

For $\gamma = 4.6$, for instance, a χ^2 fit yields that $D_0 = 9.16(36)$ and $D_1 = 3.894(77)$. If the tree level equation (5.1) were correct at the quantum level, too, then it would mean that a_4 vanishes at $\beta = D_0/D_1 = 2.35(14)$ in the deconfining phase. This would contradict the assumption that in the confining




FIG. 5. C_1^{-1} versus β for different γ 's, where C_1^{-1} is defined in Eq. (5.4). The graph shows the scaling behavior in the Coulomb phase. The lines correspond to the linear function (6.1), where D_0 and D_1 are given in Table II.

phase the lattice spacing a_4 approaches zero as β approaches ~ 3.7 for $\gamma=4.6$ (see Table I). This does not necessarily mean that the transition from the deconfining phase to the confining one is a first order transition or a crossover transition. It may be well possible that the tree-level form (5.1) receives quantum corrections in such a way that the transition is indeed of second order. Therefore, we consider possible quantum corrections to C_1^{-1} that are consistent with the scaling law in Fig. 5 and the value of β^* in the confining phase (given Table I). Since C_1^{-1} , being dimensionless, can depend only on the combination R/g_5^2 , the correction can only be a constant, i.e.,

$$C_1^{-1} \sim 2\pi R/g_5^2 + \alpha \quad \text{or} \quad C_1 \sim \frac{g_5^2}{2\pi R + \alpha g_5^2}. \quad (6.2)$$

In Table III we give the results of the fits, from which we find that the ansatz for the nonperturbative quantum correction to the coefficient of the Coulomb potential (5.4) is consistent with our data, and we conclude that

$$\alpha = 5.1 \pm 0.7, \quad (6.3)$$

where we have not included the data for $\gamma=5.0$ in Eq. (6.3), because the error for this case is very large compared with others. This indicates that the assumption that the transition from the confining to the deconfining phase is a second order transition is consistent with the data. Note that the transition

TABLE II. Fit for C_1 defined in Eq. (6.1). The fitted lines in Fig. 5 intersect with the β axis at $\beta=D_0/D_1$.

γ	D_0	D_1	$(\beta_{\min}:\beta_{\max})$	χ^2/DOF	D_0/D_1
3.6	9.48(31)	4.827(80)	(3.20 : 4.60)	0.103	1.965(97)
4.0	10.08(30)	4.603(74)	(3.40 : 4.80)	0.0957	2.19(10)
4.6	9.16(36)	3.894(77)	(4.00 : 5.40)	0.0998	2.35(14)
5.0	8.39(57)	3.47(11)	(4.20 : 5.80)	0.175	2.41(24)

TABLE III. γ independence of α .

γ	α
3.6	5.03(66)
4.0	5.05(64)
4.6	5.35(78)
5.0	5.7(11)

for small values of γ , or large values of $R\Lambda$, is of first order [6,8]. We expect that the first order transition for large values of $R\Lambda$ changes to a crossover transition, and finally to a second order transition, as we decrease the value of $R\Lambda$.

The nonperturbative correction (6.2) means that the tree level relation $\hat{g}^2 = g_5^2/2\pi R$ should be modified to

$$\hat{g}^2 = \frac{g_5^2}{2\pi R} \left(1 + \alpha \frac{g_5^2}{2\pi R} \right)^{-1}. \quad (6.4)$$

Since α is large, the correction is not small. The Coulomb phase may be of phenomenological importance, because the color degrees of freedom do not need to be always confined. The $SU(2)$ part of the standard model, for instance, could result from a higher-dimensional Yang-Mills theory in the Coulomb phase. Then an equation such as Eq. (6.4) defines the matching condition.

VII. CONCLUSION

In this paper we performed Monte Carlo simulations in a five-dimensional lattice $SU(2)$ Yang-Mills theory, where we compactified one extra dimension. We found that, as the compactification radius R decreases, the confining phase spreads more and more to the weak coupling regime, and the effective dimension of the theory gradually changes from five to four. Our data indicate that there exists a maximal radius above which the color degrees of freedom are not confined. An actual computation of the maximal radius in QCD will give an important phenomenological constraint for model building based on Kaluza-Klein theories. Our data also indicate that for fixed R/a_4 the transition from the deconfining phase to the Coulomb phase is of second order if R/a_4 is small enough.

The parameter regime we have considered in the present work corresponds to the regime in which the Kaluza-Klein idea is expected to be realized: At short distances we have the five-dimensional rotational invariance, and at long distances, the Kaluza-Klein excitations decouple so that the low-energy effective theory is a four-dimensional Yang-Mills theory. We found no indication that would contradict this picture. Moreover, the compactified five-dimensional theory, which is perturbatively nonrenormalizable, has predictive power (unless examined at very short distances), as we con-

³In the case of the phase transition measured by the Polyakov loop that extends into the fifth dimension, the change from first to second order happens at a certain value of γ [8].

clude from the scaling laws we observe. (Readers are also invited to see [17].)

The parameter regime that corresponds to deconstructing extra dimensions [14] is not the same as above [18]; the two phases are nonperturbatively separated [8,18]. In the phase for the conventional Kaluza-Klein theory, the vacuum expectation value of the Polyakov loop (which extends into the fifth dimension) is nonzero [8], while it vanishes [18] in the phase for deconstructing extra dimensions. (The phase for deconstructing extra dimensions is the one in which the layer structure in five-dimensional gauge theories can be realized [19].) Although it is not at all clear that the five-dimensional rotational invariance at short distances is recovered, it looks at the moment as if two different confining four-dimensional Yang-Mills theories could result from two different phases

(one from each) of a five-dimensional theory. The difference is purely nonperturbative. It will be very exiting to investigate this difference in more detail, especially in supersymmetric cases, where one already has analytic results, and it is shown that the five-dimensional Lorentz invariance is recovered [20].

ACKNOWLEDGMENTS

This work was supported by Grants-in-Aid for Scientific Research from the Japan Society for the Promotion of Science (JSPS) (No. 11640266, and No. 13135210). We would like to thank K.-I. Aoki, V. Bornyakov, M. Murata, H. Nakano, M. Polikarpov, G. Schierholz, H. So, T. Suzuki, and H. Terao for useful discussions.

-
- [1] Th. Kaluza, Sitzungsber. K. Preuss. Akad. Wiss. Phys. Math. K1, 966 (1921); O. Klein, Z. Phys. 37, 895 (1926).
 - [2] I. Antoniadis, Phys. Lett. B 246, 377 (1990); I. Antoniadis, C. Muñoz, and M. Os, Nucl. Phys. B397, 515 (1993).
 - [3] N. Arkani-Hamed, S. Dimopoulos, and G. Dvali, Phys. Lett. B 429, 263 (1998); Phys. Rev. D 59, 086004 (1999).
 - [4] K. Dienes, E. Dudas, and T. Gherghetta, Phys. Lett. B 436, 55 (1998); Nucl. Phys. B537, 47 (1999).
 - [5] M. Creutz, Phys. Rev. Lett. 43, 553 (1979).
 - [6] C.B. Lang, M. Pilch, and B.-S. Skagerstam, Int. J. Mod. Phys. A 3, 1423 (1988).
 - [7] H. Kawai, M. Nio, and Y. Okamoto, Prog. Theor. Phys. 88, 341 (1992); J. Nishimura, Mod. Phys. Lett. A 11, 3049 (1996).
 - [8] S. Ejiri, J. Kubo, and M. Murata, Phys. Rev. D 62, 105025 (2000).
 - [9] M. Peskin, Phys. Lett. 94B, 161 (1980).
 - [10] T.R. Taylor and G. Veneziano, Phys. Lett. B 212, 147 (1988).
 - [11] T. Kobayashi, J. Kubo, M. Mondragon, and G. Zoupanos, Nucl. Phys. B550, 99 (1999).
 - [12] D. O'Connor and C.R. Stephens, Phys. Rev. Lett. 72, 506 (1994); Int. J. Mod. Phys. A 9, 2805 (1994).
 - [13] J. Kubo, H. Terao, and G. Zoupanos, Nucl. Phys. B574, 495 (2000).
 - [14] N. Arkani-Hamed, A.G. Cohen, and H. Georgi, Phys. Rev. Lett. 86, 4757 (2001); C.T. Hill, S. Pokorski, and J. Wang, Phys. Rev. D 64, 105005 (2001).
 - [15] G. Burgers, F. Karsch, A. Nakamura, and I.O. Stamatescu, Nucl. Phys. B304, 587 (1988); T.R. Klassen, *ibid.* B533, 557 (1998); S. Ejiri, Y. Iwasaki, and K. Kanaya, Phys. Rev. D 58, 094505 (1998); J. Engels, F. Karsch, and T. Scheideler, Nucl. Phys. B564, 303 (2000).
 - [16] APE Collaboration, M. Albanese *et al.*, Phys. Lett. B 192, 163 (1987); G.S. Bali and K. Schilling, Phys. Rev. D 47, 661 (1993).
 - [17] J. Kubo and M. Nunami, hep-th/0112032.
 - [18] M. Murata, Ph.D. thesis, Kanazawa University, 2002; M. Murata and H. So (unpublished).
 - [19] Y.K. Fu and H.B. Nielsen, Nucl. Phys. B236, 167 (1984); P. Dimopoulos, K. Farakos, A. Kehagias, and G. Koutsoumbas, *ibid.* B617, 237 (2001); P. Dimopoulos, K. Farakos, and C.P. Korthals-Altes, J. High Energy Phys. 02, 005 (2001); P. Dimopoulos, K. Farakos, and S. Nicolis, hep-lat/0105014; P. Dimopoulos, K. Farakos, and G. Koutsoumbas, Phys. Rev. D 65, 074505 (2002).
 - [20] C. Csáki, J. Erlich, C. Grojean, and G.D. Kribs, Phys. Rev. D 65, 015003 (2001); C. Csáki *et al.*, *ibid.* 65, 085033 (2002).