

Table 1. Mortality rates for subtypes of cerebrovascular diseases in dialysis patients and the general population in Japan

Cerebrovascular disease	Dialysis patients		General population		Standardized mortality ratio [‡] (95% CI)
	Total No. of deaths	Crude mortality rates [†]	Crude mortality rates [†]	Age-adjusted mortality rates [†]	
Intracerebral hemorrhage	933	17.07	2.06	4.48	3.8 (3.6-4.1)
Cerebral infarction	511	9.35	3.14	7.44	1.3 (1.2-1.4)
Subarachnoid hemorrhage	147	2.69	0.99	2.00	1.3 (1.1-1.6)

[†]Per 10 000 person-years, adjusted for age distribution of dialysis patients; [‡]standardized for age; CI, confidence interval.

Table 2. Crude mortality (per 10 000 person-years, 95% confidence interval) stratified by dialysis vintage among dialysis patients

Subtype of cerebrovascular diseases	Dialysis vintage, years				
	<1y (67 154 patient- years)	1-4 (200 500 patient- years)	5-9 (137 234 patient- years)	10-14 (67 392 patient- years)	≥15 (74 194 patient- years)
	17.3 (14.3-20.7)	16.1 (14.4-18.0)	17.8 (15.6-20.2)	18.3 (15.2-21.8)	16.6 (13.8-19.8)
Intracerebral hemorrhage					
Cerebral infarction	8.3 (6.3-10.8)	10.4 (9.1-11.9)	9.9 (8.3-11.7)	8.0 (6.0-10.5)	7.0 (5.2-9.2)
Subarachnoid hemorrhage	3.0 (1.8-4.6)	2.2 (1.6-3.0)	2.8 (2.0-3.9)	3.4 (2.2-5.1)	2.4 (1.4-3.8)

Figure 1.

