

Table 1 Number of deaths as a result of nephrotic syndrome and crude mortality rates by sex and age groups

	Range (calendar years)							
	1995-1999		2000-2004		2005-2009		2010-2014	
	Number of deaths	Mortality rate (95% CI)	Number of deaths	Mortality rate (95% CI)	Number of deaths	Mortality rate (95% CI)	Number of deaths	Mortality rate (95% CI)
Males								
Age 0-19 years	18	0.03 (0.02-0.04)	9	0.01 (0.01-0.03)	4	0.01 (0.002-0.02)	6	0.01 (0.004-0.02)
20-39	11	0.01 (0.01-0.02)	18	0.02 (0.01-0.03)	9	0.01 (0.01-0.02)	9	0.01 (0.01-0.02)
40-59	99	0.11 (0.09-0.14)	86	0.10 (0.08-0.12)	69	0.08 (0.06-0.10)	57	0.07 (0.05-0.09)
60-79	522	1.00 (0.92-1.09)	514	0.85 (0.78-0.93)	548	0.81 (0.75-0.88)	531	0.71 (0.65-0.77)
≥80	452	6.48 (5.90-7.11)	477	5.57 (5.08-6.09)	580	4.95 (4.55-5.37)	853	5.59 (5.22-5.98)
Overall	1102	0.36 (0.34-0.38)	1104	0.36 (0.34-0.38)	1210	0.39 (0.37-0.42)	1456	0.47 (0.45-0.50)
Females								
Age 0-19 years	20	0.03 (0.02-0.05)	7	0.01 (0.01-0.02)	5	0.01 (0.003-0.02)	5	0.01 (0.003-0.02)
20-39	6	0.01 (0.003-0.02)	5	0.01 (0.002-0.01)	7	0.01 (0.003-0.02)	2	0.003 (0.0003-0.01)
40-59	53	0.06 (0.04-0.08)	53	0.06 (0.05-0.08)	39	0.05 (0.03-0.06)	12	0.02 (0.01-0.03)
60-79	446	0.71 (0.64-0.78)	377	0.54 (0.48-0.59)	351	0.46 (0.41-0.51)	310	0.37 (0.33-0.41)
≥80	687	4.83 (4.48-5.21)	735	4.02 (3.73-4.32)	885	3.72 (3.48-3.97)	1068	3.65 (3.43-3.87)
Overall	1212	0.38 (0.36-0.40)	1177	0.37 (0.35-0.39)	1287	0.40 (0.38-0.42)	1397	0.43 (0.41-0.45)

Per 100 000 person-years.

Table 2 Annual percentage change (APC) and 95% confidence intervals (95% CI) for age-standardised mortality rates (ASR), as determined by joinpoint regression analysis for the years 1995-2014

	1995 ASR	2014 ASR	1st period trend		
			Period	APC (95% CI)	P-value
Men	0.26	0.18	1995-2014	1.9% (2.3% to 1.4%)	<0.01
Women	0.18	0.10	1995-2014	3.5% (4.1% to 2.9%)	<0.01

ASR per 100 000 person-years, according to the World Standard Population.

Table 3 Estimated years of life lost (YLL) and average years of life lost (AYLL) owing to nephrotic syndrome or malignant neoplasm by sex and age in 1995, 2000, 2005, and 2010 using the constant endpoint method

	YLL				AYLL			
	1995	2000	2005	2010	1995	2000	2005	2010
Men								
Nephrotic syndrome	785	486	339	484	19.6	13.1	14.8	12.7
Malignant neoplasm	504 719	461 429	396 438	333 019	9.7	9.5	8.7	8.1
Women								
Nephrotic syndrome	389	241	280	153	16.9	11.0	14.0	13.9
Malignant neoplasm	372 799	357 641	321 067	286 466	12.0	11.6	10.8	10.2