# Structural and functional analysis of the eukaryotic ribosomal stalk complex by site-directed mutagenesis

Kentaro Baba

Doctoral Program in Life and Food Sciences Graduate School of Science and Technology Niigata University

# CONTENTS

ABSTRACT	3
ABBREVIATIONS	4
INTRODUCTION	5-8
MATERIALS AND METHODS	9-11
Plasmid Construction, Protein Expression, and Purification	9
In Vitro Reconstitution of the Stalk Complex	10
Ribosomal 50 S Core and Hybrid 50S Particles	10
Quantitative Analysis of P1-P2 Heterodimer Incorporated into the Ribosome	11
Ribosome Functional Assays	11
RESULTS	12-19
	12 15
Binding sites for P1-P2 at the C-terminal helices of P0	12-13
Binding sites for P1-P2 at the C-terminal helices of P0 Contributions of individual copies of P1 and P2 to ribosome function	12-13
Binding sites for P1-P2 at the C-terminal helices of P0 Contributions of individual copies of P1 and P2 to ribosome function Role of the conserved YPT motif between the two spine helices I and II	12-13 15-17 17-19
Binding sites for P1-P2 at the C-terminal helices of P0 Contributions of individual copies of P1 and P2 to ribosome function Role of the conserved YPT motif between the two spine helices I and II FIGURES	12-13 15-17 17-19 <b>20-27</b>
Binding sites for P1-P2 at the C-terminal helices of P0 Contributions of individual copies of P1 and P2 to ribosome function Role of the conserved YPT motif between the two spine helices I and II FIGURES TABLE	12-13 15-17 17-19 20-27 28
Binding sites for P1-P2 at the C-terminal helices of P0 Contributions of individual copies of P1 and P2 to ribosome function Role of the conserved YPT motif between the two spine helices I and II FIGURES TABLE DISCUSSION	12-13 15-17 17-19 20-27 28 29-33
Binding sites for P1-P2 at the C-terminal helices of P0 Contributions of individual copies of P1 and P2 to ribosome function Role of the conserved YPT motif between the two spine helices I and II FIGURES TABLE DISCUSSION REFERENCES	12-13 15-17 17-19 20-27 28 29-33 34-40

#### ABSTRACT

In animal ribosomes, two stalk proteins P1 and P2 form a heterodimer, and the two dimers, with the anchor protein P0, constitute a pentameric complex crucial for recruitment of translational GTPase factors to the ribosome. To investigate the functional contribution of each copy of the stalk proteins, we constructed P0 mutants, in which one of the two C-terminal helices, namely helix I (N-terminal side) or helix II (C-terminal side) were unable to bind the P1-P2 dimer. We also constructed "one-C-terminal domain (CTD) stalk dimers", P1-P2 $_{\Delta C}$  and P1 $_{\Delta C}$ -P2, composed of intact P1/P2 monomer and a CTD-truncated partner. Through combinations of P0 and P1-P2 variants, various complexes were reconstituted and their function tested in eEF-2-dependent GTPase and eEF-1 $\alpha$ /eEF-2-dependent polyphenylalanine synthesis assays in vitro. Double/single-CTD dimers bound to helix I showed higher activity than that bound to helix II. Despite very low polypeptide synthetic activity by a single one-CTD dimer, its binding to both helices considerably increased activity, suggesting that two stalk dimers cooperate, particularly in polypeptide synthesis. This promotion of activity by two stalk dimers was lost upon mutation of the conserved YPT sequence connecting the two helices of P0, suggesting a role for this sequence in cooperativity of two stalk dimers.

## **ABBREVIATIONS**

 $A_{260}$ ; absorbance at 260 nm aP1; archaeal ribosomal protein P1 cpm; count per minute Cryo-EM; Cryo-electron microscopy eEF-1α; eukaryotic elongation factor 1α eEF-2; eukaryotic elongation factor 2 EF-G; elongation factor G EF-Tu; elongation factor Tu eL12; eukaryotic ribosomal protein L12 mRNA; messenger RNA NMR; nuclear magnetic resonance PAGE; polyacrylamide gel electrophoresis PCR; polymerase chain reaction rpm; rotation per minute rRNA; ribosomal RNA SDS; sodium dodecyl sulfate WT; wild-type

#### INTRODUCTION

The dynamic process of mRNA translation is facilitated by interactions of the ribosome with several translational GTPase factors (1-3). The ribosomal large subunits contain a characteristic protein complex in all organisms, termed the stalk complex, responsible for the recruitment of GTPases and GTP hydrolysis (4-9). The stalk complex is composed of multiple copies of a flexible acidic protein, or so-called stalk protein, together with the anchor protein or stalk base. The stalk protein, a key component of this functional unit, is composed of an Nterminal domain (NTD) that participates in dimerization (6, 10-14), a C-terminal domain (CTD) responsible for direct binding to GTPases (9, 15-18) and a flexible hinge region connecting the N- and C-domains (19). Despite sharing a common functionality, structures and assembly modes of stalk complexes are markedly different between the three domains of life.

In bacteria, the stalk protein L12 forms a homodimer by hydrophobic interaction between the N-terminal 30 residue region, which forms antiparallel V-shaped  $\alpha$ 1- $\alpha$ 2 hairpins. Subsequently, two or three L12-L12 homodimers bind via their NTDs to the C-terminal spine helix of the anchor protein L10 (6). The CTD of bacterial L12 is a globular structure that is composed of 70 amino acids (20). Binding of the CTD to the GTPase factors EF-G, EF-Tu, and IF-2 has been demonstrated by analyses of NMR chemical shifts (15), cross-linking (16), cryo-EM (17), and X-ray crystallography (18). In contrast, the eukaryotic stalk proteins P1/P2 and the archaeal counterpart aP1 have a large N-terminal dimerization domain composed of about 70 amino acids, which form a globular  $\alpha 1 \cdot \alpha 4$  structure (12, 14) and a small CTD composed of about 20 amino acids, which is responsible for binding to GTPases (9). Archaeal aP1 forms a homodimer and three dimers bind to the C-terminal spine helices of the anchor protein aPO (12) while eukaryotic P1 and P2 form a heterodimer (13, 14, 21-23) and two P1-P2 heterodimers bind to the C-terminal helices of P0 (24-27). Eukaryotic P1/P2 and archaeal aP1 are related closely to each other, but not to bacterial L12, with respect to amino acid sequence (28, 29). It has been therefore suggested that eukaryotic/archaeal stalk protein might not be linked evolutionally to bacterial L12.

Given the essential role of the stalk protein complex in ribosome function, the distinct variations of structure and organization of the complexes in eukaryotes, archaea, and bacteria appear to reflect their characteristic functional modes within the different domains of life. The eukaryotic stalk complex is of particular interest. It is composed of three components P0, P1 and P2, which share a common CTD (30). In lower eukaryotes (e.g., Saccharomyces cerevisiae), P1 and P2 have evolved into four different components P1 $\alpha$ , P1 $\beta$ , P2 $\alpha$ , and P2 $\beta$ (31, 32). The most prominent characteristic of eukaryotic stalks including those from S. cerevisiae is that they form a P1-P2 heterodimer, whose structure is asymmetrical (13, 14), unlike the archaeal aP1-aP1 homodimer (12). Starting from this asymmetric feature of the P1-P2 structure, Lee et al. recently proposed that two heterodimers assemble on the C-terminal spine helices of P0 in a P2-P1:P1-P2 mode, in which P1 faces itself via hydrophobic interactions (14). In this model, the two neighbouring P1 components form a cavity that accommodates a loop region containing the conserved YPT motif between the two helices. The conserved YPT sequence, therefore, might be relevant to cooperativity in binding between P0 and P1-P2. In addition, it has been shown by cross-linking experiments that the locations of the CTDs of P1 and P2 in the S. cerevisiae ribosome are different, suggesting distinct roles for P1 and P2 on the ribosome (33). These characteristics of eukaryotic stalk complex are likely to be related to the sophisticated regulation of the stalk assembly and translation mechanism, although there remain issues to be clarified.

The present study focuses on the functional contribution of each copy of the stalk proteins, P1 and P2. P0 mutants lacking the ability to bind to either one of

the two P1-P2 dimers were constructed. The stalk complexes were reconstituted *in vitro* with the P0 mutants, together with one-CTD stalk dimer lacking the CTD of either P1 or P2, and assayed for their activity in GTPase assays and in polypeptide synthesis dependent on eukaryotic elongation factors by using a hybrid ribosome system (34). We could show not only the activity of individual copies of the stalks, but also functional cooperativity between the two stalk dimers. We also detected a novel functional contribution of the conserved YPT motif between two spine helices of P0, to which P1-P2 dimers bind individually, suggesting that the YPT region is involved in the functional cooperativity of the two stalk dimers.

#### MATERIALS AND METHODS

#### **Plasmid Construction, Protein Expression, and Purification**

The cDNAs for *B. mori* ribosomal protein full-length P0 and the truncation mutant lacking the C-terminal 55 amino acids (P0 $\Delta C$ ) were amplified by PCR and inserted into the *NcoI* and *Bam*HI sites of pET15b (Novagen). Double mutations F215Q/V222Q in helix I (P0mutI) and I241Q/L248Q in helix II (P0mutII) were introduced by a PCR-based site-directed mutagenesis method (35). The same methodology was used for replacement of the 231-YPTI-234 sequence in the loop region between helices I and II with 231-GGGG-234 (P0mutM), as well as for deletion of this sequence ( $PO_{\Delta M}$ ). The coding sequences for *B. mori* ribosomal proteins P1, P2, and eL12 (a eukaryotic homologue of prokaryotic L11) were cloned into pET28c or pET3a (Novagen), as described previously (25). The truncation mutants lacking the C-terminal 52 amino acids of P1 (P1 $_{\Delta C}$ ) and lacking 50 amino acids of P2 (P2 $\Delta c$ ) were constructed as described previously (36). The nucleotide sequences of the constructed plasmids were confirmed by DNA sequencing analyses. Expression and purification of each ribosomal protein was performed as described previously (25). The purity of all proteins was checked by SDS-PAGE.

#### In Vitro Reconstitution of the Stalk Complex

The stalk complexes were reconstituted by mixing isolated P0, P1, and P2, or alternative individual mutants, in the presence of 7 M urea at a molar ratio of 1:3:3, as described previously (22). An rRNA fragment containing residues 1841– 1939 of rat 28 S rRNA (corresponding to residues 1043–1112 of *E. coli* 23 S rRNA) was synthesized *in vitro* using template DNA and SP6 RNA polymerase (25). The P0-P1-P2-rRNA complex was formed by adding 3-fold amounts of the rRNA fragment to 100 pmol of the P0-P1-P2 complex. Individual complexes were analysed by 6 % polyacrylamide gel electrophoresis (22). The gel was stained with Coomassie Brilliant Blue.

#### **Ribosomal 50S Core and Hybrid 50S Particles**

Ribosomal 50S subunits were prepared from the L11-deficient *E. coli* strain AM68 (37), as previously described (34). The 50S core particles deficient both in L10 and L7/L12 were prepared by extraction of the L11-deficient 50S subunits in a solution containing 50 % ethanol and 0.5 mM NH<sub>4</sub>Cl at 0 °C, as previously described (34). The *B. mori-E. coli* hybrid 50S particle was formed by mixing the *E. coli* 50S core with the reconstituted stalk complex P0-P1-P2 and eL12 (25).

#### Quantitative Analysis of P1-P2 Heterodimer Incorporated into the Ribosome

*B. mori* ribosomal protein P2 was radiolabeled by phosphorylation using  $[\gamma^{-32}P]$ ATP and casein kinase II (New England Biolabs), as described previously (25). The stalk complex and its variants were reconstituted with the labeled [<sup>32</sup>P]P2, as described above. The labeled stalk complexes (90 pmol) were incubated with 50S core particles (30 pmol) supplemented with eL12 (90 pmol) in a solution (20 µl) that contained 20 mM Tris-HCl (pH 7.6), 5 mM MgCl<sub>2</sub>, 50 mM NH<sub>4</sub>Cl, and 5 mM 2-mercaptoethanol at 37 °C for 10 min. The mixture was layered onto a 10-28 % sucrose gradient and centrifuged at 40,000 rpm at 4 °C for 3 h in a Hitachi S52ST rotor, as described previously (25). The fraction that contained the 50S particles was collected, and the amount of associated P1-P2 dimer was estimated on the basis of the amount of radioactivity, considering the specific radioactivity of [<sup>32</sup>P]P2 to be 840 cpm/pmol.

#### **Ribosome Functional Assays**

Eukaryotic eEF-1 $\alpha$  and eEF-2 were isolated from pig liver, as previously described (38). The activity of the hybrid ribosome with respect to eEF-2-dependent GTPase activity and eEF-1 $\alpha$  /eEF-2-dependent polyphenylalanine synthesis was assayed as described previously (25 and 34).

#### RESULTS

# Binding sites for P1-P2 at the C-terminal helices of P0

Based on previous data on the archaeal stalk complex, we assumed that the hydrophobic residues F215, G218, V222, and L225 in helix I, and I241, G244, L248, and I251 in helix II of *B. moli* P0, which are located at positions -5, -2, +2, and +5 in the conserved 2-fold symmetric distribution (Fig. 1 and 2A), are involved in binding of P1-P2 heterodimer in each of the helicies I and II of P0. We constructed the P0 mutants P0mutI and P0mutII, which contained double mutations at positions -5 and +2, i.e., F215Q/V222Q and I241Q/L248Q, respectively. These P0 mutants were expressed in *E. coli* cells and purified, as described in "MATERIALS AND METHODS" (Fig. 3). Each of the P0 mutants was mixed with excess amounts of P1-P2 heterodimer in vitro, and complex formation was analysed by native gel electrophoresis (Fig. 4A). In contrast to the failure of isolated P0 to run into the gel (Fig. 4A, lane 1), and to the smearing pattern of isolated P1 obtained because of its tendency to aggregate (Fig. 4A, lane 2) (39), the P0-P1-P2 stalk complex produced a distinct band (lane 5), whose mobility was clearly lower than that of isolated P2 (lane 3) and P1-P2 heterodimer (lane 4), as described previously (22). The complexes were also formed with

P0mutI (lane 6) and P0mutII (lane 7) mutants. The mobility of the complexes with the mutant P0 was slightly lower than that of the wild-type P0 (lane 5). It is noteworthy that, compared with the gel pattern of the wild-type complex sample, there remained larger amounts of free P1-P2 dimer with high gel mobility in samples containing the mutant P0 (lanes 6 and 7), and that the band corresponding to the P0mutI-P1-P2 complex (lane 6) was thinner than that for the P0mutII-P1-P2 complex (lane 7). It has been reported that binding of P0 to an rRNA fragment covering the P0 binding site stabilizes a variant of the P0-P1-P2 complex (25). We therefore performed the same gel electrophoresis after adding the rRNA fragment. As shown in Fig. 4B, the electrophoretic mobility of the wild-type P0-P1-P2 complex (Fig. 4B, lane 1) became slightly higher upon addition of the RNA (lane 3). The complexes containing the P0 mutants also migrated faster after addition of the RNA (Fig. 4B, lanes 4 and lanes 5) when compared with the same samples without the RNA (Fig. 4A, lanes 6 and lanes 7, respectively), suggesting that the RNA fragment bound to the P0mutI-P1-P2 and P0mutII-P1-P2 complexes. The thin band produced by the P0mutI-P1-P2 complex observed in Fig. 4A (lane 6) became distinct upon RNA addition (Fig. 4B, lane 4), suggesting that the RNA binding to P0mutI stabilizes the complex structure of P0mutI-P1-P2. It is again noticeable that even after addition of the RNA, larger

amounts of free P1-P2 were observed in samples including the P0 mutants (lanes 4 and 5), compared with the wild-type P0 sample (lane 3). These results suggest that the P0mutI and P0mutII mutants may bind only a single P1-P2 heterodimer.

To determine stoichiometry of the complexes P0mutI-P1-P2 and P0mutII-P1-P2, we used the reconstituted radiolabeled P1-[32P]P2 heterodimer in the in *vitro* assembly of the stalk complexes, as described previously (25). The labeled stalk complex P0-P1-<sup>[32</sup>P]P2 formed by mixing of P0 and excess amounts of P1-[<sup>32</sup>P]P2 was substituted for the *E. coli* stalk complex L10-L12 on the 50S subunit. This eukaryote-bacterium hybrid form, a so-called the "hybrid ribosome" has been successfully used for studies on assembly and function of eukaryotic and archaeal stalk complexes (34, 40). In the present study, the P1-[<sup>32</sup>P]P2 heterodimer bound to P0 or its mutants and assembled onto the *E. coli* 50S core were separated by sucrose gradient centrifugation (Fig. 5). Radioactivity in the 50S fraction was measured, and copy numbers of P1-P2 heterodimer were estimated on the basis of the amounts of 50S particles and bound P1-P2 that were estimated from the A<sub>260</sub> and radioactivity values, respectively. The results, summarized in Table 1, clearly indicate that both P0mutI and P0mutII mutants have the ability to bind only a single copy of [<sup>32</sup>P]P2, implying a single P1-P2 heterodimer binds to

individual P0 mutants. Therefore, the amino acid substitutions introduced into helix I and helix II (Fig. 1) appear to disrupt P1-P2 binding to individual helices.

#### Contributions of individual copies of P1 and P2 to ribosome function

It has long been a question as to why eukaryotic ribosomes have two sets of heterodimers composed of P1 and P2, which share a common CTD, in particular with regards to their functional contribution. In the present study, we used P0mutI and P0mutII mutants, which allow us to investigate the functional contributions of individual P1-P2 dimers at the different locations. To focus on the roles of P1 and P2, we used a  $P0_{\Delta C}$  mutant lacking the CTD of P0, a sequence shared by P1 and P2. Moreover, we constructed and used the one-CTD stalk dimers P1-P2<sub> $\Delta C$ </sub> and P1<sub> $\Delta C$ </sub>-P2, in which the CTD of either P1 or P2 is truncated (see MATERIALS AND METHODS). The stalk complexes were reconstituted by various combinations of P0 and P1-P2 variants (Fig. 6), and incorporated onto the *E. coli* 50S core. The resultant hybrid ribosomes were assayed for their activity in eEF-2-dependent GTPase and eEF-1a/eEF-2-dependent polyphenylalanine synthesis assays (Fig 7).

The hybrid ribosomes containing P0mutII<sub> $\Delta C$ </sub>-P1-P2 (carrying the single P1-P2 dimer bound to helix I) have low but significant levels of activity in GTPase assays (73 % of P0<sub> $\Delta C$ </sub>-P1-P2) and polyphenylalanine synthesis (38 % of P0<sub> $\Delta C$ </sub>-P1-P2), whereas these activities of the ribosomes containing P0mutI<sub> $\Delta C$ </sub>-P1-P2 (carrying a single P1-P2 dimer bound to helix II) were only 35 % and 12 % that of the  $P0_{\Delta C}$  stalk complex carrying double P1-P2 dimers, respectively (Fig. 7A). The results indicate that P1-P2 dimer bound to helix I makes a larger contribution to the factor dependent activities than the dimer bound to helix II. The same characteristics were also observed when one-CTD stalk dimers P1- $P2_{\Delta C}$  and  $P1_{\Delta C}$ -P2 were used, instead of the P1-P2 dimer, although the activities of one-head stalk dimers were lower than those of P1-P2 dimer (Fig. 7B). It is noticeable that the P1-CTD stalk dimer (P1-P2<sub>AC</sub>) has slightly higher activities than P2-CTD dimer (P1 $_{\Delta C}$ -P2).

The activities of the hybrid ribosomes containing  $P0_{\Delta C}$ -P1-P2<sub> $\Delta C$ </sub> and  $P0_{\Delta C}$ -P1<sub> $\Delta C$ </sub>-P2 (carrying double one-CTD dimers bound to both helicies I and II) were also assayed (Fig. 7C). The level of GTPase activity of each double one-CTD stalk complex was approximately the sum of the activities of single one-CTD stalk dimers bound to helices I and II (compare Fig. 7C with B). In contrast, the level of polypeptide synthesis of each double one-CTD stalk complex was obviously higher than the sum of activities of individual ones (Fig. 7C and B). The results suggest that the stalk bound to helix I cooperates with that bound to helix II, particularly in polypeptide synthesis.

It should be noted that the CTD of P0, which is shared by P1 and P2, also contributes to ribosome function (41). To examine its contribution to eEF-2dependent GTPase and eEF-1 $\alpha$ /eEF-2-dependent polyphenylalanine synthesis, we also performed functional assays with intact P0 carrying the CTD, instead of the P0<sub>AC</sub> mutant, and compared the data (Fig. 7D) with that in Fig 7A. Although all of P0-P1-P2, P0mutI-P1-P2, and P0mutII-P1-P2 showed slightly higher activity (Fig. 7D) than individual variants lacking the CTD of P0 (Fig. 7A), the effect of mutations at helicies I (P0mutI) and II (P0mutII) was roughly comparable with the effects observed with P0mutI<sub>AC</sub> and P0mutII<sub>AC</sub>. It is therefore likely that the CTD of P0 exerts its basal functionality, but both P1-P2 dimers do not depend on the presence of the CTD of P0.

#### Role of the conserved YPT motif between the two spine helices I and II

There is a conserved sequence motif, Thy-Pro-Thr, within a loop region between helices I and II of eukaryotic P0 and archaeal aP0. To examine the functional role of this region, we constructed the variants P0mutM, in which the 231-YPTI-234 sequence was changed to 231-GGGG-234, and PO<sub>AM</sub>, in which the same sequence was deleted (Fig. 8). These P0 variants were expressed in *E. coli* cells and purified (Fig. 3). The ability of P0mutM and P0<sub> $\Delta M$ </sub> to bind P1-P2 heterodimer was analysed by native gel electrophoresis in the presence of the rRNA fragment covering the P0 binding site (Fig. 8A). A distinct band was observed for the stalk-rRNA complex containing P0mutM (Fig. 8A, lane 2) and the intensity of the band was comparable with that of the wild-type complex (lane 1 and 5). The result suggests that the YPT motif does not have an essential role in P1-P2 binding. To confirm this suggestion, we introduced additional mutations into P0mutM, i.e., the P1-P2 binding site of either helix I (lane 3) or helix II (lane 4) was disrupted in P0mutM. The results clearly indicate that even after introduction of these mutations, individual P0 mutants still retain the ability to bind P1-P2 dimer (lanes 3 and 4), suggesting that the mutation of the 231-YPTI-234 sequence produced no detectable effect on individual binding of P1-P2 to helices I and II. To examine the functional effect of mutation of the 231-YPTI-234 sequence, the stalk complexes containing P0mutM were assembled into the 50S core and assayed for polyphenylalanine synthesis activity (Fig. 8B). Unexpectedly, the activity of the complex containing P0mutM was significantly reduced (33 % of

P0-P1-P2), although both helices I and II of this mutant retain their P1-P2 binding ability (Fig. 8A). A similar result was also obtained with the truncation mutant P0<sub> $\Delta M$ </sub> (Fig. 8A, lane 6 and B). It is noteworthy that the levels of activity of the complexes containing P0mutM and P0<sub> $\Delta M$ </sub> were comparable with the complex containing P0mutII (36 % of P0-P1-P2), which can bind a single P1-P2 dimer to helix I.

#### FIGURES



# Figure 1. Sequence comparison of the stalk binding sites on the C-terminal helices of eukaryotic P0/archaeal aP0.

Amino acid sequences around the C-terminal helices I and II of eukaryotic P0 (Bmo, Bombix mori; Hsa, Homo sapiens; Pfa, Plasmodium falciparum; Ath, Arabidopsis thaliana; Ddi, Dictyostelium discoideum; and Sce, Saccharomyces cerevisiae) and the helices I, II, and III of archaeal aPO (Sac, Sulfolobus acidocaldarius; Pae, Pyrobaculum aerophilum; Mva, Methanococcus vannielii; Hma, Haloarcula marismortui; Tac, Thermoplasma acidophilum; and Pho, Pyrococcus horikoshii) were compared. Helical regions defined by X-ray crystallography of PhoP0 (12) are represented as cylinders. Hydrophobic residues involved in binding of PhoP1 homodimers are located in 2-fold symmetric distribution at  $\pm 2$  and  $\pm 5$ . These residues are colored orange for residues at -5 and -2 and green for residues at +2 and +5. Mutations introduced are indicated above the sequences. The conserved YPT sequence located between helices I and II is colored red. Mutations introduced in this study are indicated above the sequences and in the schematic models of the eukaryotic stalk complex. In this model, PO is represented as three tandemly linked boxes and a circle (corresponding to the N-terminal domain, helix I, helix II, and the CTD of P0, respectively), and P1 and P2 are represented in red and blue, respectively. The mutated helices are marked with a green X. Introduction of the mutations in the YPT sequence is represented by M surrounded by a green circle in these models.



Figure 2. Model of binding of the stalk dimer to a C-terminal helix of aP0.

(A) Close-up view of the archaeal aP1-aP1 homodimer (white/pink) bound to the spine helix I (cyan) of aP0. The hydrophobic residues important for binding are shown as green and orange sticks (PDB accession code 3A1Y) (12). (B) Superposition of human P1-P2 heterodimer (P1, red; P2, dark blue, PDB accession code 2LBF) (14) and the archaeal aP1-aP1 homodimer (white), with aP0-helix I (cyan). Hydrophobic residues of P1 (I10 and L14) and P2 (V4 and L8), which are located at close proximity to the conserved hydrophobic residues of aP0 (L217, A220, A224 and L227), are represented as sticks.



# Figure 3. SDS-PAGE pattern of isolated protein samples.

Individual samples (2 µg of each): P0 (lane 1); P0mutI (lane 2); P0mutII (lane 3); P0mutM (lane 4); P0<sub> $\Delta$ M</sub> (lane 5); P0<sub> $\Delta$ C</sub> (lane 6); P0mutI<sub> $\Delta$ C</sub> (lane 7); P0mutII<sub> $\Delta$ C</sub> (lane 8); P1 (lane 9); P1<sub> $\Delta$ C</sub> (lane 10); P2 (lane 11); P2<sub> $\Delta$ C</sub> (lane 12); eL12 (lane 13) were subjected to SDS-PAGE on a 16.5 % acrylamide separation gel. The gel was stained with Coomassie Brilliant Blue.



# Figure 4. Formation of stalk complexes with P0 carrying mutations in helices I and II.

(A) The complexes were reconstituted by incubation of P0 (lane 5), P0mutI (lane 6) and P0mutII (lane 7) with excess amounts of P1-P2 dimer. Each complex sample including 100 pmol of P0 (or its mutants) was subjected to native gel electrophoresis, as described in "MATERIALS AND METHODS". On the same gel, isolated P0 (100 pmol, lane 1), P1 (300 pmol, lane 2), P2 (300 pmol, lane 3) and P1-P2 dimer (300 pmol, lane 4) were also loaded. The gel was stained with Coomassie Brilliant Blue. (B) The complexes were reconstituted by incubation of P0 (lanes 1, 3), P0mutI (lane 4), and P0mutII (lane 5) with P1-P2 dimer under the same conditions as (A). Each complex including 100 pmol of P0 (or its mutant) was incubated in the presence of the rRNA fragment (300 pmol) for the P0 binding site (lanes 3-5) or in its absence (lane 1), and subjected to native gel electrophoresis under the same conditions as (A). On the same gel, isolated P0 (100 pmol) mixed with the rRNA fragment was loaded (lane 2). The gel was stained with Coomassie Brilliant Blue. Schematic models of the stalk complexes are shown below individual lanes.



# Figure 5. Quantitative analysis of stalk dimers bound to P0 carrying mutations in helices I and II in the 50S particle.

The 50S core particles (30 pmol) were mixed with excess amounts of the P0-P1-P2 complexes (90 pmol), which are composed of P1-[<sup>32</sup>P]P2 dimers (840 cpm/pmol) and P0, and eL12 (90 pmol) (A). The hybrid 50S particles were also reconstituted using P0mutI (B), and P0mutII (C), instead of P0. The hybrid 50S samples were separated by sucrose gradient centrifugation, as described in "MATERIALS AND METHODS". The amount of P2 incorporated to the 50S core was estimated by radioactivity of P2 and the absorbance at 260 nm of the 50S core.



### Figure 6. Formation of variants of the stalk complex.

The complexes were reconstituted by incubation in combinations as follows:  $PO_{\Delta C}$  with P1-P2 (lane 1),  $PO_{\Delta C}$  with P1-P2<sub> $\Delta C$ </sub> (lane 2),  $PO_{\Delta C}$  with P1<sub> $\Delta C$ </sub>-P2 (lane 3),  $PO_{\Delta C}$  with P1<sub> $\Delta C$ </sub>-P2<sub> $\Delta C$ </sub> (lane 4), P0mutI<sub> $\Delta C$ </sub> with P1-P2 (lane 5), P0mutI<sub> $\Delta C$ </sub> with P1-P2<sub> $\Delta C$ </sub> (lane 6), P0mutI<sub> $\Delta C$ </sub> with P1<sub> $\Delta C$ </sub>-P2 (lane 7), P0mutII<sub> $\Delta C$ </sub> with P1-P2 (lane 8), P0mutII<sub> $\Delta C$ </sub> with P1-P2<sub> $\Delta C$ </sub> (lane 9) and P0mutII<sub> $\Delta C$ </sub> with P1<sub> $\Delta C$ </sub>-P2 (lane 10). Each sample containing 100 pmol of P0 variant was mixed with 300 pmol of the rRNA fragment covering the P0 binding site, and separated by native gel electrophoresis, as described in "MATERIALS AND METHODS". The gel was stained with Coomassie Brilliant Blue. Schematic models of the stalk complexes are shown below individual lanes.



Figure 7. Contributions of individual copies of P1 and P2 to eEF-2-dependent GTPase activity and polyphenylalanine synthesis.

The stalk complexes used, as represented in the schematic diagrams for the stalk complexes below the bars, were reconstituted in various combinations with the P0 and P1-P2 variants: (A) P0<sub>AC</sub>, P0mutI<sub>AC</sub>, or P0mutII<sub>AC</sub>, together with P1-P2, or P1<sub>AC</sub>-P2<sub>AC</sub>; (B) P0<sub>AC</sub>, P0mutI<sub>AC</sub>, or P0mutII<sub>AC</sub>, together with P1-P2, P1-P2<sub>AC</sub>, or P1<sub>AC</sub>-P2; (D) P0, P0mutI, P0mutII, or P0<sub>AC</sub> together with P1-P2. For the GTPase acivity, *E. coli* 50S core particles (2.5 pmol) were preincubated with each reconstituted P0-P1-P2 variant (10 pmol) and 10 pmol eL12, and then assayed for eukaryotic eEF-2-dependent GTPase activity (white bar). The 100 % activity corresponds to 80.5 pmol (A, B and C) or 87.0 pmol (D) of [<sup>32</sup>P]GTP hydrolysis per min. For the polypeptide synthetic activity, the 50S core particles (10 pmol) were preincubated with 20 pmol of the same P0-P1-P2 samples and eL12, and then assayed for polyphenylalanine synthesis dependent on eukaryotic eEF-1 $\alpha$  and eEF-2 (black bar). The 100 % activity corresponds to 16.6 pmol (A, B and C) or 20.5 pmol (D) of [<sup>14</sup>C] phenylalanine polymerized for 10 min.



Figure 8. Role of the conserved YPT motif between the two spine helices I and II.

(A) The stalk complexes, as represented in the schematic diagrams of the stalk complexes below the bars, were reconstituted in combinations of P1-P2 dimers with P0 (lane 1 and 5), P0mutM (lane 2), P0mutI/mutM (lane 3), P0mutII/mutM (lane 4) and P0<sub> $\Delta$ M</sub> (lane 6). Each complex sample containing 100 pmol of P0 (or its variant) with 300 pmol of the rRNA fragment for the P0 binding region was subjected to native gel electrophoresis, as described in "MATERIALS AND METHODS". The gel was stained with Coomassie Brilliant Blue. (B) The 50S core particles (10 pmol) were preincubated with the stalk complex sample P0-P1-P2, P0mutI-P1-P2, P0mutI-P1-P2, P0mutI-P1-P2, P0mutI-P1-P2, represented to reach and eL12 (20 pmol), and then assayed for polyphenylalanine synthesis dependent on eukaryotic eEF-1 $\alpha$  and eEF-2. The 100 % activity corresponds to 21.0 pmol of [<sup>14</sup>C] phenylalanine polymerized for 10 min.

## TABLE

Table 1. Copy numbers for P2 bound to the 50S core particle in the presence of different P0 variants.

P0 type	Copies of P2
WT	2.22 ± 0.14
mutl	$1.04 \pm 0.14$
mutll	$1.14 \pm 0.09$

The copy number of labeled [<sup>32</sup>P]P2 (840 cpm/ pmol) per 50S particle was calculated from absorbance measurements at 260 nm and the radioactivity of the 50S fractions separated by sucrose gradient centrifugation shown in Fig. 5, as described in "MATERIALS AND METHODS".

#### DISCUSSION

The animal ribosomal stalk is composed of P0 and two copies of P1-P2 heterodimers, which individually bind to two tandem helices (I and II) of P0. In the present study, we identified hydrophobic amino acid residues F215/V222 in helix I and I241/L248 in helix II, as being crucial for individual P1-P2 binding to the helices. The hydrophobicity of these residues is conserved among eukaryotes and archaea. In the crystal structure of the archaeal aP0-aP1-aP1 stalk complex, it has been shown that four hydrophobic residues at positions -5, -2, +2, and +5 in a 2-fold symmetric distribution within each helix interact with  $\alpha 1/\alpha 1$  of the aP1aP1 homodimer. The residues at -5 and -2 make contact with  $\alpha 1$  in one aP1 molecule of the homodimer and the residues at +2 and +5 make contact with  $\alpha 1$ ' of the other aP1 molecule. We infer that the mode of binding between eukaryotic P0 and P1-P2 is similar to that of archaeal aP0 and aP1 homodimer on the basis of the following facts; (i) the 2-fold symmetric distribution of the hydrophobic residues is conserved from archaea to eukaryotes (Fig. 1); (ii) the residues F215/V222 in helix I and I241/L248 in helix II at positions corresponding to -5 and +2 are crucially important for P1-P2 binding as revealed in the present study; (iii) it has been reported that the  $\alpha 1/\alpha 1$  of eukaryotic P1 and P2 are important for their binding to P0 as well as for dimerization (36, 42). It is therefore likely that the four conserved hydrophobic residues in each the helix of eukaryotic P0 participate in binding of the P1-P2 heterodimer. To explore this possibility, we attempted to fit the crystal structural data for helix I of archaeal aP0 (12) to the NMR structural data for human P1-P2 dimer ( $\alpha$ 1 to  $\alpha$ 3) (14). In one such model (Fig. 2B), the four conserved hydrophobic residues of aP0 are able to bind to hydrophobic residues of the  $\alpha$ 1/ $\alpha$ 1' helices of P1 and P2 (Fig. 2B), i.e., the residues at positions -5 and -2 of P0 can make contact with  $\alpha$ 1 of P2. Meanwhile, the residues at positions +2 and +5 can make contact with  $\alpha$ 1' of P1, assuming that the two P1-P2 dimers assemble on the spine helices of P0 in a P2-P1:P1-P2 mode, as proposed by Lee et al. (14). This putative model, however, contrasts with models proposed previously, in which P1 plays an anchoring role for P2 (43, 44).

The resultant mutants P0mutI (F215Q/V222Q) and P0mutII (I241Q/L248Q) actually bind only a single P1-P2 heterodimer (Fig. 5). Using the P0mutI and P0mutII mutants, together with the one-CTD stalk heterodimers lacking the CTD of either P1 or P2, we investigated the functional contributions of individual copies of the P1/P2 stalk. Lower but considerable levels of eEF-2-dependent GTPase activity were shown by the P1-P2 dimer (Fig. 7A) or one-CTD dimers (Fig. 7B) bound only to helix I, compared with the activity in the control stalk complex carrying double P1-P2 dimers. Further reduced levels of GTPase activity were also detected in the same dimer samples bound only to helix II. In contrast, further lowering of activity in  $eEF-1\alpha/eEF-2$ -dependent polyphenylalanine synthesis assays was observed in the dimer samples bound only to helix I, and a background level of activity was present in dimer samples bound only to helix II. The results suggest that the stalk dimer bound to helix II might make little contribution to polypeptide synthesis. However, when the P1-P2 dimer or one-CTD dimers bind to both helices I and II, high levels of polypeptide synthetic activity were observed (Fig. 7A and C). It is therefore more likely that the P1-P2 stalk dimer bound to helix II works cooperatively with the stalk dimer bound to helix I, particularly in polypeptide synthesis. This may relate to the fact that two elongation factors eEF-1 $\alpha$  and eEF-2 are required for the peptide polymerization reaction, and that collaboration of two stalk dimers may be required for alternate action of the two elongation factors for efficient polypeptide synthesis. The present experimental system provides straightforward evidence for functional collaboration of the two stalk dimers bound to helices I and II.

The different nature of the CTDs of P1 and P2 within the yeast ribosome has been revealed by previous cross-linking experiments (33); the CTD of P2 shows much higher reactivity towards several other ribosomal components than the CTD of P1. The present study shows that the P1-head dimer, composed of P1 and  $P2_{\Delta C}$  lacking the CTD, shows slightly but reproducibly higher activity than the P2-head dimer, composed of P2 and P1\_{\Delta C}, with respect to polypeptide synthesis as well as eEF-2-dependent GTPase activity (Fig. 7B and C). Our results indicate that both P1 and P2 contribute to ribosomal activity dependent on eEF-1 $\alpha$  and eEF-2, and that the degree or quality of contributions seems to be slightly different between P1 and P2. Although it is hard to explain a correlation between the present functional data and previous cross-linking data at present, further detailed investigations with both approaches should provide evidence for differences in the functional nature of P1 and P2.

There is a loop region containing a conserved sequence motif, Thy-Pro-Thr, between helices I and II in eukaryotic P0 and archaeal aP0. Little information is available on the functional role of this region although Lee et al. (14) proposed an interesting hypothesis where they suggest that when two P1-P2 dimers bind individually to helices I and II, the  $\alpha$ 3 regions of the P1 molecules face each other by hydrophobic interaction and make a hydrophobic cavity, into which the YPT loop can fit. The present study demonstrated that a change of amino acid sequence or deletion of the YPT loop gave no detectable effect on binding of P1-P2 to both the helices of P0 (Fig. 8A), but caused a marked effect on polypeptide synthesis (Fig. 8B). The present experimental data indicate that the YPT loop is involved in function of the stalk complex, rather than the assembly of the stalk complex. It is however unlikely that the YPT loop plays a role in direct contact with translation factors, because of its probable location within a hydrophobic cavity (12, 14). Considering the fact that the level of reduced activity of the YPT loop-defective mutants is comparable with that of P0mutII lacking the P1-P2 dimer bound to helix II, it seems likely that the disruption of the YPT loop causes loss in function of the stalk dimer bound to helix II by changing the angle or orientation of helix II with respect to helix I. Because the stalk dimer bound to helix II apparently collaborates with the dimer bound to helix I as discussed above, the YPT loop would seem to play a role in the cooperation of the two stalk dimers.

#### REFERENCES

- 1. Kaziro,Y. (1978) The role of guanosine 5'-triphosphate in polypeptide chain elongation. *Biochim. Biophys. Acta*, **505**, 95-127.
- 2. Bourne,H.R., Sanders,D.A. and McCormick,F. (1991) The GTPase superfamily: conserved structure and molecular mechanism. *Nature*, **349**, 117-127.
- Liljas, A. (2004) Structural Aspects of Protein Synthesis, (World Scientific Publishing Co., Singapore), pp99-158.
- Wahl,M.C. and Möller,W. (2002) Structure and function of the acidic ribosomal stalk proteins. *Curr. Protein Pept. Sci.*, **3**, 93-106.
- Gonzalo, P. and Reboud, J.P. (2003) The puzzling lateral flexible stalk of the ribosome. *Biol. Cell.*, 95, 179-193. Review.
- 6. Diaconu,M., Kothe,U., Schlünzen,F., Fischer, N., Harms,J.M., Tonevitsky,A.G., Stark,H., Rodnina,M.V. and Wahl,M.C. (2005) Structural basis for the function of the ribosomal L7/12 stalk in factor binding and GTPase activation. *Cell*, **121**, 991-1004.
- Savelsbergh, A., Mohr, D., Kothe, U., Wintermeyer, W. and Rodnina, M.V. (2005) Control of phosphate release from elongation factor G by ribosomal protein L7/12. *EMBO J.* 24, 4316-4323.
- Mandava,C.S., Peisker,K., Ederth,J., Kumar,R., Ge,X., Szaflarski,W. and Sanyal,S. (2012) Bacterial ribosome requires multiple L12 dimers for efficient initiation and elongation of protein synthesis involving IF2 and EF-G. *Nucleic Acids Res.*, 40, 2054-2064.

- 9. Nomura,N., Honda,T., Baba,K., Naganuma,T., Tanzawa,T., Arisaka,F., Noda,M., Uchiyama,S., Tanaka,I., Yao,M. and Uchiumi,T. (2012) Archaeal ribosomal stalk protein interacts with translation factors in a nucleotideindependent manner via its conserved C terminus. *Proc. Natl. Acad. Sci. USA*, 109, 3748-3753.
- Wahl,M.C., Bourenkov,G.P., Bartunik,H.D. and Huber,R. (2000) Flexibility, conformational diversity and two dimerization modes in complexes of ribosomal protein L12. *EMBO J.*, **19**, 174-86.
- Briceño, V., Camargo, H., Remacha, M., Santos, C. and Ballesta, J.P. (2009) Structural and functional characterization of the amino terminal domain of the yeast ribosomal stalk P1 and P2 proteins. *Int. J. Biochem. Cell Biol.*, 41, 1315-1322.
- Naganuma, T., Nomura, N., Yao, M., Mochizuki, M., Uchiumi, T. and Tanaka, I.
   (2010) Structural basis for translation factor recruitment to the eukaryotic/archaeal ribosomes. J. Biol. Chem., 285, 4747-4756.
- Tchórzewski, M., Boldyreff, B., Issinger, O.G. and Grankowski, N. (2000) Analysis of the protein-protein interactions between the human acidic ribosomal P-proteins: evaluation by the two hybrid system. *Int. J. Biochem. Cell Biol.*, **32**, 737-746.
- Lee,K.M., Yu,C.W., Chiu,T.Y., Sze,K.H., Shaw,P.C. and Wong,K.B. (2012) Solution structure of the dimerization domain of the eukaryotic stalk P1/P2 complex reveals the structural organization of eukaryotic stalk complex. *Nucleic Acids Res.*, 40, 3172-82.

- Helgstrand, M., Mandava, C.S., Mulder, F.A., Liljas, A., Sanyal, S. and Akke, M. (2007) The ribosomal stalk binds to translation factors IF2, EF-Tu, EF-G and RF3 via a conserved region of the L12 C-terminal domain. *J. Mol. Biol.*, 365, 468-479.
- Nechifor,R. and Wilson,K.S. (2007) Crosslinking of translation factor EF-G to proteins of the bacterial ribosome before and after translocation. *J. Mol. Biol.*, 368, 1412-1425.
- Harms,J.M., Wilson,D.N., Schluenzen,F., Connell,S.R., Stachelhaus,T., Zaborowska,Z., Spahn,C.M. and Fucini,P. (2008) Translational regulation via L11: molecular switches on the ribosome turned on and off by thiostrepton and micrococcin. *Mol. Cell*, **30**, 26-38.
- Gao,Y.G., Selmer,M., Dunham,C.M., Weixlbaumer,A., Kelley,A.C. and Ramakrishnan,V. (2009) The structure of the ribosome with elongation factor G trapped in the posttranslocational state. *Science*, **326**, 694-699.
- Dey,D., Oleinikov,A.V. and Traut,R.R. (1995) The hinge region of *Escherichia* coli ribosomal protein L7/L12 is required for factor binding and GTP hydrolysis. *Biochimie*, 77, 925-930.
- Leijonmarck, M. and Liljas, A. (1987) Structure of the C-terminal domain of the ribosomal protein L7/L12 from *Escherichia coli* at 1.7 A. *J. Mol. Biol.*, 195, 555-79.
- Zurdo, J., González, C., Sanz, J.M., Rico, M., Remacha, M. and Ballesta, J.P. (2000) Structural differences between Saccharomyces cerevisiae ribosomal stalk proteins P1 and P2 support their functional diversity. *Biochemistry*, 39, 8935-8943.

- 22. Shimizu,T., Nakagaki,M., Nishi,Y., Kobayashi,Y., Hachimori,A. and Uchiumi,T. (2002) Interaction among silkworm ribosomal proteins P1, P2 and P0 required for functional protein binding to the GTPase-associated domain of 28S rRNA. *Nucleic Acids Res.*, **30**, 2620-2627.
- Tchórzewski, M., Krokowski, D., Boguszewska, A., Liljas, A. and Grankowski, N.
   (2003) Structural characterization of yeast acidic ribosomal P proteins forming the P1A-P2B heterocomplex. *Biochemistry*, **142**, 3399-408.
- 24. Pérez-Fernández, J., Remacha, M. and Ballesta, J.P. (2005) The acidic protein binding site is partially hidden in the free Saccharomyces cerevisiae ribosomal stalk protein P0. *Biochemistry*, 44, 5532-5540.
- 25. Hagiya,A., Naganuma,T., Maki,Y., Ohta,J., Tohkairin,Y., Shimizu,T., Nomura,T., Hachimori,A. and Uchiumi,T. (2005) A mode of assembly of P0, P1, and P2 proteins at the GTPase-associated center in animal ribosome: *in vitro* analyses with P0 truncation mutants. *J. Biol. Chem.*, **280**, 39193-39199.
- 26. Krokowski, D., Boguszewska, A., Abramczyk, D., Liljas, A., Tchórzewski, M. and Grankowski, N. (2006) Yeast ribosomal P0 protein has two separate binding sites for P1/P2 proteins. *Mol. Microbiol.*, **60**, 386-400.
- 27. Smulski, C.R., Longhi, S.A., Ayub, M.J., Edreira, M.M., Simonetti, L.,
  Gómez, K.A., Basile, J.N., Chaloin, O., Hoebeke, J. and Levin, M.J. (2011)
  Interaction map of the Trypanosoma cruzi ribosomal P protein complex
  (stalk) and the elongation factor 2. *J. Mol. Recognit.*, 24, 359-370.
- 28. Shimmin,L.C., Ramirez,C., Matheson,A.T. and Dennis,P.P. (1989) Sequence alignment and evolutionary comparison of the L10 equivalent and L12

equivalent ribosomal proteins from archaebacteria, eubacteria, and eucaryotes. J. Mol. Evol., 29, 448-462.

- Grela, P., Bernadó, P., Svergun, D., Kwiatowski, J., Abramczyk, D., Grankowski, N., and Tchórzewski, M. (2008) Structural relationships among the ribosomal stalk proteins from the three domains of life. *J. Mol. Evol.*, 67, 154-167.
- 30. Rich, B.E. and Steitz, J.A. (1987) Human acidic ribosomal phosphoproteins P0,
  P1, and P2: analysis of cDNA clones, *in vitro* synthesis, and assembly. *Mol. Cell Biol.* 7, 4065-4074.
- 31. Guarinos, E., Remacha, M. and Ballesta, J.P. (2001) Asymmetric interactions between the acidic P1 and P2 proteins in the Saccharomyces cerevisiae ribosomal stalk. J. Biol. Chem., 276, 32474-32479.
- 32. Tchórzewski, M. (2002) The acidic ribosomal P proteins. Int. J. Biochem. Cell Biol., 34, 911-915. Review.
- 33. Qiu,D., Parada,P., Marcos,A.G., Cárdenas,D., Remacha,M. and Ballesta,J.P.
  (2006) Different roles of P1 and P2 Saccharomyces cerevisiae ribosomal stalk proteins revealed by cross-linking. *Mol Microbiol.*, 62, 1191-1202.
- 34. Uchiumi,T., Honma,S., Nomura,T., Dabbs,E.R. and Hachimori,A. (2002) Translation elongation by a hybrid ribosome in which proteins at the GTPase center of the *Escherichia coli* ribosome are replaced with rat counterparts. *J. Biol. Chem.*, **277**, 3857-3862.
- 35. Kunkel,T.A., Bebenek,K. and McClary,J. (1991) Efficient site-directed mutagenesis using uracil-containing DNA. *Methods Enzymol.*, 204, 125-139.

- 36. Naganuma,T., Shiogama,K. and Uchiumi,T. (2007) The N-terminal regions of eukaryotic acidic phosphoproteins P1 and P2 are crucial for heterodimerization and assembly into the ribosomal GTPase-associated center. *Genes Cells.* 12, 501-510.
- 37. Dabbs, E.R. (1979) Selection for *Escherichia coli* mutants with proteins missing from the ribosome. *J. Bacteriol.*, **140**, 734-737.
- Iwasaki,K. and Kaziro,Y. (1979) Polypeptide chain elongation factors from pig liver. *Methods Enzymol.*, 60, 657-676.
- 39. Grela,P., Sawa-Makarska,J., Gordiyenko,Y., Robinson,C.V., Grankowski,N. and Tchórzewski,M. (2008) Structural properties of the human acidic ribosomal P proteins forming the P1-P2 heterocomplex. *J. Biochem.*, 143, 169-177.
- 40. Nomura, T., Nakao, K., Maki, Y., Naganuma, T., Takashima, T., Tanaka, I., Kimura, M., Hachimori, A. and Uchiumi, T. (2006) *In vitro* reconstitution of the GTPase-associated centre of the archaebacterial ribosome: the functional features observed in a hybrid form with *Escherichia coli* 50S subunits. *Biochem. J.*, **396**, 565-571.
- 41. Remacha, M., Jimenez-Diaz, A., Bermejo, B., Rodriguez-Gabriel, M.A., Guarinos, E. and Ballesta, J.P. (1995) Ribosomal acidic phosphoproteins P1 and P2 are not required for cell viability but regulate the pattern of protein expression in Saccharomyces cerevisiae. *Mol. Cell. Biol.*, 15, 4754-4762.
- 42. Camargo,H., Nusspaumer,G., Abia,D., Briceño,V., Remacha,M., and Ballesta,J.P. (2011) The amino terminal end determines the stability and

assembling capacity of eukaryotic ribosomal stalk proteins P1 and P2. Nucleic Acids Res., **39**, 3735-3743.

- 43. Zurdo, J., Parada, P., van den Berg, A., Nusspaumer, G., Jimenez-Diaz, A., Remacha, M., and Ballesta, J.P. (2000) Assembly of Saccharomyces cerevisiae ribosomal stalk: binding of P1 proteins is required for the interaction of P2 proteins. *Biochemistry*, **39**, 8929-8934.
- 44. Gonzalo,P., Lavergne,J.P. and Reboud,J.P. (2001) Pivotal role of the P1 Nterminal domain in the assembly of the mammalian ribosomal stalk and in the proteosynthetic activity. *J. Biol. Chem.*, **276**, 19762-19769.

## ACKNOWLEDGEMENTS

I would like to express my sincere gratitude to my supervisors Dr. Toshio Uchiumi for all of his grate supports, excellent suggestions and hard work during the course of this dissertation. This thesis would never be completed without his support. And I would like to thank all former and present members of the Uchiumi lab for companionship and support.