

Scrotal Skin Temperature Affects the Activity of the Hypothalamic Magnocellular Neurons in the Rat

Takao AKAISHI

Division of Body Fluid Physiology, Course for Biological Functions and Medical Control, Graduate School of Medical and Dental Sciences, Niigata University, Niigata, Japan

Received October 16 2002; accepted November 22 2002

Summary. Single unit activities in the hypothalamic supraoptic nucleus (SON) were antidromically identified as a magnocellular neurosecretory cell in male Wistar rats under urethane anesthesia (1.2 g/kg, b. w.). Forty-two unit of these activities were categorized as 30 presumed-vasopressin (p-AVP) and 12 presumed oxytocin producing (p-OXT) cells, and were examined in their responsiveness to thermal stimuli topically applied to the scrotal skin. The stimulated area was kept at 30°C throughout the experiment. Warming (to 43°C) of the scrotal skin caused an inhibition in the discharge rate of p-AVP cells. Cooling (to 15°C) had no effect. In the p-OXT cell, neither warming nor cooling thermal stimuli had an effect on their firing activity. Neither thermal stimulus had any effect on the heart rate and mean blood pressure. These results indicate that information from the scrotal thermoreceptors affects the neuroendocrinal activity of the hypothalamic magnocellular neuroendocrine system and that AVP may take part in the thermoregulatory mechanism to maintain body temperature in the rat.

Key words—supraoptic neuron, vasopressin, oxytocin, scrotal skin temperature, rat.

INTRODUCTION

In the rat, thermal stimulation of the scrotal skin influences neuronal activity in the hypothalamus⁸, septum⁴, and thalamus¹⁵. The magnocellular neuron in the hypothalamic paraventricular nucleus was suggested to be responsive to thermal stimulation of the preoptic and anterior hypothalamus^{9,10}. However,

little is known about the afferent connection from the peripheral thermoreceptors to the hypothalamic magnocellular neuron. This study was therefore designed to elucidate the afferent neural connections and its characteristic features between the thermoreceptors on the scrotal skin and the magnocellular neuron of the hypothalamic supraoptic nucleus (SON) in the rat.

METHODS AND MATERIALS

A total of 23 Wistar male rats housed in a controlled environment (20–23°C) with a 14:10 hr light-dark cycle (lights on at 7 a.m.) were used in the present series of experiments. Food and water were provided *ad libitum*. The rats, weighing between 250 to 360 g, were anesthetized with urethane given *i. p.* at a dose of 1.2 g/kg. The rectal temperature was maintained at 37.5°C throughout the experiment by means of a warming pad (ATB-1100, Nihon Khoden co.). The electrocardiogram (ECG) was lead from the left leg and right arm. In some rats, the right femoral artery was cannulated to measure blood pressure (BP). The rats were mounted in a stereotaxic instrument (SR-6, Narishige) in a prone position. Craniotomy was done in the parietal area, and the sagittal sinus was ligated and removed. A side-by-side bipolar stimulating electrode, constructed from stainless steel wire 200 μm thick (F-2D, Narishige), was inserted into the neurohypophysis at the midline. Recordings were made with glass micropipettes constructed from a Pyrex

Correspondence: Takao Akaishi, Division of Body Fluid Physiology, Course for Biological Functions and Medical Control, Graduate School of Medical and Dental Sciences, Niigata University, Asahimachidori 1, Niigata 951-8510, Japan

Abbreviations—BP, blood pressure, ECG, electrocardiogram; HR, heart rate; p-AVP, presumed vasopressin; p-OXT, presumed oxytocin; SON, supraoptic nucleus; SST, scrotal skin temperature.




Fig. 1 A representative rate meter recording of the response of a supraoptic p-AVP neuron to warming (43°C) stimuli of the dorsal scrotal skins (A). The upper smoothed superimposed three responses were obtained from another p-AVP neuron during repetitive warming stimuli (B).

tube (GDO-1.5, Narishige) filled with a 2% (wt/vol) solution of Pontamine sky blue dye in a 0.5 M sodium acetate buffer. The direct current resistance of the recording electrode was 6–11 M Ω . The SON was systematically explored while isolated rectangular pulses of 0.5 ms duration, at a 100–1500 μA intensity, were applied at 0.81 Hz through the stimulation electrode inserted into the neurohypophysis. Penetrations of recording electrodes were confined to an area 1.5–2.0 mm lateral to the midline, 7.3–8.0 mm rostral from the lamda, and 0.2–0.7 mm dorsal from the ventral hypothalamic surface according to the atlas by Albe-Fessard *et al.*³⁾. Conventional procedures were followed for amplification and display of the extracellular action potentials according to our previous study¹⁾. The antidromic identification of discharge activities of the p-AVP producing cells was followed by criteria proposed by Poulain and Wakerley¹²⁾.

For thermal stimulation of the scrotal skin, the dorsal region of the scrotal skin was clipped and unfolded. An attached surface (about 10.0 mm²) of the thermode was placed on the midline of the dorsal portion so that it adhered to the skin of the scrotum. Skin temperature was measured by a thermocouple implanted inside the thermode that was in contact with the skin (BTC-201, Unique Medical).

The scrotal skin temperature (SST) was kept at 30°C before and after stimulation, and SST was set and maintained at 43 or 15°C for 5 min during the examination of responses of a particular SON neur-

on. Warming stimuli were always tested first, with cooling ones following the warming ones. In some responsive neurons, stimuli were repeatedly applied. The control and reaired mean discharge rate of a particular neuron was determined from 3 min before until 3 min after the stimulus.

All measurements, rectal temperature, SST, BP, heart rate (HR), and single neuronal activity were introduced to the MacLab system (MacLab/400, Bioresearch Center) via a regular or conventional high gain amplifier for microelectrode recording (MEZ-8201, Nihon Kohden Kogyo), and were analyzed.

ANOVA and Duncan's method were applied for analysis of the effect of stimulus on the SON neuronal activity. For determination of the mean response of the single unit activity, a simple smoothing method (normalizing points: 101) was applied.

All aspects of the present experiments were conducted in conformity with the guiding principles for the care and use of animals approved by the council of the Japan Physiological Society (2001).

RESULTS

The extracellular unit activity of a total of 51 neurons was recorded from the SON in the male rat ($N=23$). Among them, 30 neurons exhibited a phasic pattern of the spontaneous discharge. An important




Fig. 2 Each neuronal response (thinner lines) and mean response (thicker curve line) to warming (A) and cooling stimuli (B) of the scrotal skin in all p-AVP neurons examined. Using a moving average method, each rate meter recording (thinner lines) are smoothed and superimposed. Thicker straight line shows the scrotal skin temperature (SST, °C).

Table 1. Effects of cooling and warming of the scrotal skin on the burst activity of the supraoptic p-AVP neuron

Parameter	Cooling (15°C) N=16		Warming (43°C) N=28	
	Control	Stimulus	Control	Stimulus
Number of burst (Burst/min)	4.9±0.8*	4.7±1.0	5.5±0.7	3.3±0.5**
Interval of burst (sec)	18.5±3.4	17.7±3.0	14.5±2.6	19.8±2.6
Firiny rate in burst (spikes/sec)	7.5±0.6	7.4±0.5	7.4±0.4	7.5±0.5

*; mean±SEM, **; P<0.05.

Table 2. The mean blood pressure (MBP) and heart rate (HR) before (3 min), during (5 min) and after (3 min) the warming or cooling stimulus on the scrotal skin in 4 rats

Stimulus		Before	During	After
Warming (N****=16)	MBP*	69.3±3.7***	69.5±3.7	68.9±3.9
	HR**	310.8±7.1	310.0±7.5	312.1±7.8
Cooling (N=16)	MBP	70.8±3.7	70.6±3.6	70.1±3.7
	HR	305.1±7.7	304.8±7.8	305.0±7.6

*, mmHg; **, beats/min; ***, mean±SEM; ****, trial number.

criterion for the classification as a phasic type was that the firing discharge activity was interrupted by distinct periods of complete silence, which began and terminated abruptly during observations. Other SON 12 neurons had a firing pattern with a constant discharge rate, and were designated as a continuous type in this study. Thus, they satisfied the necessary conditions proposed by Poulain and Wakerley¹²⁾. According their study, the former type of neurons were presumed “vasopressin (p-AVP) neurons”, the latter “oxytocin (p-OXT) neurons”. These two types of unit activities were recorded well within the nuclear boundary of the SON. Nine SON cells presented such difficulty in determining their obvious firing pattern because of their lower discharge rate that they were excluded from further examination.

A representative example of reversible changes in the neuronal activity of p-AVP cells when the SST was changed from 30°C to 43°C is shown in Fig. 1. The control firing discharge rates of p-AVP cells before warming stimuli were 5.0 ± 0.7 (N=28) spikes/sec. The warming temperature in SST significantly inhibited the firing discharge rate (3.6 ± 0.6 Hz, $P < 0.001$) of these examined cells. The control firing discharge rates of p-AVP cells before cooling stimuli were 3.6 ± 0.5 Hz spikes/sec (N=16). The cooling in SST, on the other hand, had induced insignificant changes in the discharge rate in p-AVP cells (3.9 ± 0.7 spikes/sec, *N. S.*) during cooling. The discharge rates after stimuli were 4.8 ± 0.8 (warming) and 4.0 ± 0.7 Hz (cooling), respectively. Each response in all p-AVP neurons examined is shown in Fig 2.

The burst activities for all p-AVP cells examined were further analyzed before and during application of the stimulus. The mean number, interval and firing rate of the burst are shown in Table 1. The warming stimulus significantly decreased the number of bursts generated per minute ($P < 0.05$). The two other parameters in warming and cooling were not affected.

In the p-OXT cells examined, no significant change in their discharge rate could be detected during either warming (1.2 ± 0.2 Hz, N=11) or cooling (1.5 ± 0.3 Hz, N=11) stimuli when compared with that during the control period (1.3 ± 0.2 and 1.5 ± 0.2 Hz, respectively).

Mean blood pressure (MBP) and heart rate (HR) were measured continually from 4 rats and the effects of thermal stimuli on the scrotal skin were tested. Neither MBP nor HR, showed significant changes during either warming or cooling stimuli as shown in Table 2.

DISCUSSION

Thermoreceptor afferent inputs derived from the scrotal skin to the hypothalamus have been reported in a number of past studies^{6,8,14,15,17)}. However, to date there has been no direct evidence showing that the magnocellular neuroendocrine system receives the thermal influences from the peripheral organs, including the skin. This study is the first to show that the supraoptic p-AVP neurons were inhibited by the warming stimuli applied to the scrotal skin without variation of the body temperature.

Local hypothalamic cooling causes a diuresis in the goat¹⁶⁾ and monkey⁵⁾, and warming causes antidiuresis in the dog⁹⁾. These reports suggest that the hypothalamic temperature, in other words, the body core temperature, affects AVP release. In male rats, similar results were reported by Matsumura *et al*¹⁰⁾, namely, that the paraventricular neurosecretory neurons with a phasic firing pattern, p-AVP cells, were accelerated by warming of the preoptic and anterior hypothalamus.

Considering these past studies collectively suggests that an increase or decrease in body core temperature respectively induces an acceleration or inhibition of the AVP release for thermoregulatory function. As is generally known in thermoregulation for rela-

tively small mammals such as the rat and mouse, the role of the peripheral skin thermoreceptors should play a more important role in the regulation of the body temperature than the central hypothalamic and visceral thermoreceptors. The different heat capacity of the body and ratio of the body surface/weight may cause the body to install a different thermoregulation mechanism. Indeed, AVP is well known as a very potent vasoconstrictor in the peripheral vessel, followed by endothelium *in vitro*¹³). Therefore, in the rat, an increased or decreased release of AVP may respectively induce a vasoconstriction or vasodilatation of the skin, including the scrotum, as an initial stage of the thermoregulation. In fact, at the level of the hypothalamus, AVP was suggested to work as an antipyretic in the case of a fever¹¹). Further studies are required to clarify the relationship between the altered plasma AVP concentration and the initial stage of the thermoregulation.

Another possibility for explanation of the paradox is that the applied thermal stimuli were noxious. However, this is unlikely because the temperature stimuli applied in this study were 15 and 43°C. These temperatures have not been considered to be noxious⁴), and are within the range used in previous reports^{7,10}). Furthermore, in our previous study²), we revealed that the paraventricular p-OXT cell responds to a pinching stimulus applied to the skin. These were always accompanied with a transient increase in MBP and HR. In this study, however, p-OXT cells were not at all responsive to the thermal stimuli, and neither BP nor HR were affected during stimuli. This may directly indicate that the applied thermal stimulus was not noxious.

Another proposal that the changed cardiovascular factors may indirectly exert influence on the firing activity of the p-AVP cells examined is also unlikely because the MBP and HR during the thermal stimuli did not significantly vary.

The time course that could be detected in the mean response of the p-AVP cell was not entirely stimulus dependent. It is worth noting that the delay of the peak response was about 2–3 min. This would depend upon characteristics of the thermal receptors and/or the expanse of the stimulated area on the scrotal skin rather than the central mechanism. In fact, it is generally well known that the discharge activity of the thermal afferent of the skin is a so-called “rapidly adapting” one. The combined total of the activated afferent signals may make a final characteristic response in the p-AVP cell of the hypothalamus. Nevertheless, the possibility that it partially depends upon the central neuronal mechanism can not be neglected in this study.

In conclusion, the present study is first to demonstrate that the SST (scrotal skin temperature) affects the firing or releasing activity of the p-AVP but not that of the p-OXT cell in the SON. This result strongly suggests the AVP may take part in the initial stage of the thermoregulatory mechanism in the male rat.

REFERENCES

- 1) Akaishi T, Homma S: Estrogen modifies the baroreceptor-induced responses of supraoptic vasopressin neurons in the ovariectomized rat. *Neurosc Lett* **219**: 33–36, 1996.
- 2) Akaishi T, Robbins A, Sakuma Y, Sato Y: Neural inputs from the uterus to the paraventricular magnocellular neurons in the rat. *Neurosc Lett* **84**: 57–62, 1988.
- 3) Albe-Fessard D, Stutinsk F, Libouban S: Atlas stéréotaxique de diencephale du rat blanc. Editions du centre national de la recherche scientifique, Paris, 1966.
- 4) Disturnal J, Veale WL, Pittman QL: The ventral septal area: Electrophysiological evidence for putative arginine vasopressin projections onto thermoresponsive neurons. *Neuroscience* **19**: 795–802, 1986.
- 5) Hayward JN, Baker MA: Diuretic and thermoregulatory responses to preoptic cooling in the monkey. *Am J Physiol* **214**: 843–850, 1968.
- 6) Hellon RF, Misra NK: Neurones in the ventrobasal complex of the rat thalamus responding to scrotal skin temperature change. *J Physiology (Lond)* **232**: 389–399, 1973.
- 7) Hellon RF, Hensel H, Schafer K: Thermal receptors in the scrotum of the rat. *J Physiology (Lond)* **248**: 340–357, 1975.
- 8) Ishikawa Y, Nakayama T, Kanosue K, Matsumura K: Activation of central warm-sensitive neurons and the tail vasomotor response in rats during brain and scrotal thermal stimulation. *Pflüger Arch* **400**: 222–227, 1984.
- 9) Keil R, Gerstberger R, Simon E: Hypothalamic thermal stimulation moderates vasopressin release in hyperosmotic stimulated rabbits. *Am J Physiol (Reg Int Comp Physiol)* **267**: R1089–1097, 1994.
- 10) Matsumura K, Nakayama T, Ishikawa Y: Effects of preoptic thermal stimulation on electrical activities of neurosecretory cells in paraventricular and periventricular nuclei of the hypothalamus. *Brain Res* **289**: 330–333, 1983.
- 11) Pittman QJ, Martin S, Neumann I, Wilkinson MF, Takahashi Y: Central actions of arginine vasopressin to inhibit endotoxin and cytokine action. In: Saito T, Kurokawa K, Yoshida S (ed) *Neurohypophysin: Recent Progress of Vasopressin*. Elsevier Science Amsterdam, Lausanne, New York, Shannon, Tokyo 1995, p 321–329.
- 12) Poulain DA, Wakerley JB: Electrophysiology of

- hypothalamic magnocellular neurones secreting oxytocin and vasopressin. *Neuroscience* **7**: 773-808, 1982.
- 13) Reid IA, Schwartz J: Role of vasopressin in the control of blood pressure. In: Martini L, Ganong WF: *Frontiers in Neuroendocrinology*, Raven Press, New York 1984, p177-197.
 - 14) Shibata M, Hori T, Kiyohara T, Nakashima T: Convergence of skin and hypothalamic temperature signals on the sulcal prefrontal cortex in the rat. *Brain Res* **443**: 37-46, 1988.
 - 15) Schingnitz G: Neuronal responses in the rat's thalamus to scrotal heating. *Exp Brain Res* **43**: 419-421, 1981.
 - 16) Simon-Opperman C, Jessen C: Antidiuretic responses to thermal stimulation of hypophalamus and spinal cord in the conscious goat. *Pfuger Arch* **368**: 33-37, 1977.
 - 17) Thornhill QLJ: Thermoresponsiveness of posterior hypothalamic (PH) neurons of rats to scrotal and abdominal thermal stimulation. *Brain Res* **80**: 80-87, 1978.